

JC02 Rec'd PCT/PTO 01 APR 2002

TRANSMITTAL LETTER OF THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		Attorney Docket No. 0512-1030
		U.S. Application No. 10/089612
INTERNATIONAL APPLN. NO. PCT/FR00/02596	INTERNATIONAL FILING DATE 19 SEPTEMBER 2000	PRIORITY DATE CLAIMED 1 OCTOBER 1999
TITLE OF INVENTION: PLANT SEED ENDOSPERM-SPECIFIC PROMOTERS		
APPLICANT(S) FOR DE/EO/US: JEAN-FRANCOIS BONELLO, PETER ROGOWSKY, PASCUAL PEREZ		
Applicant herewith submits to the United States Designated Elected Office (DO/EO/US) the following items and other information:		
<ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c)(2)) <ol style="list-style-type: none"> a. <input checked="" type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau) b. <input type="checkbox"/> has been communicated by the International Bureau. See attached PCT/IB/308. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371 (c)(2)) <ol style="list-style-type: none"> a. <input checked="" type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made, however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 		
Items 11 to 20 below concern document(s) or information included:		
<ol style="list-style-type: none"> 11. <input checked="" type="checkbox"/> Information Disclosure Statement (IDS) w/PTO-1449 - <input checked="" type="checkbox"/> Copy of IDS citations 12. <input type="checkbox"/> Assignment Papers (cover sheet & document(s)) 13. <input checked="" type="checkbox"/> A FIRST Preliminary Amendment. 14. <input type="checkbox"/> A SECOND or SUBSEQUENT Preliminary Amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application (35 U.S.C. 154(d)(4)). 20. <input checked="" type="checkbox"/> Other items or information: <u>International Search Report, PCT/IB/304, PCT/IPEA/409, Abstract of the Disclosure on a Separate Sheet, Application Data Sheet</u> 		

JC10 Rec'd PCT/PTO 01 APR 2002

U.S. APPLICATION NO. 10/089612		INTERNATIONAL APPLN. NO. PCT/FR00/02596		ATTORNEY DOCKET NO. 0512-1030	
21. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1)-(5):					
Neither international preliminary examination fee nor international search fee paid to USPTO and international Search Report not prepared by the EPO or JPO \$1040.00					
International preliminary examination fee not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00					
International preliminary examination fee not paid to USPTO but International search fee paid to USPTO \$740.00					
International preliminary examination fee paid to USPTO but all claims did not satisfy provision of PCT Article 33 (1)-(4) \$710.00					
International preliminary examination fee paid to USPTO and all claims satisfied provision of PCT Article 33 (1)-(4) \$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT				\$ 890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20- <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e))				\$ 130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	29 - 20 =	9	X \$18.00	\$ 162.00	
Independent Claims	20 - 3 =	17	X \$84.00	\$ 1,428.00	
MULTIPLE DEPEND CLAIM(S) (if applicable)			+ \$280.00	\$	
TOTAL OF ABOVE CALCULATION -				\$ 2,480.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				+	
SUBTOTAL =				\$ 2,480.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492Z(f)).				\$	
TOTAL NATIONAL FEE =				\$ 2,480.00	
Fee for recording the enclosed assigned (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property +				\$	
TOTAL FEES ENCLOSED -				\$ 2,480.00	
				Amount to be refunded:	\$
				Charged:	\$
<input checked="" type="checkbox"/> A Check in the amount of \$2,480.00 to cover all fees is attached.					
<input type="checkbox"/> The Commissioner is hereby authorized to charge indicated fees and credit any overpayments to Deposit account No. 25-0120 in the name of Young & Thompson, as described below. A duplicate copy of this sheet is enclosed.					
<input checked="" type="checkbox"/> The Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 25-0120 for any additional fee required under 37 C.F.R. §§ 1.16 or 1.17.					
SEND ALL CORRESPONDENCE TO:					
745 South 23 rd Street					
Arlington, VA 22202					
Telephone (703) 521-2297					
Y&T Customer No. 000466					
		SIGNATURE <u>Benoit Castel</u>			
		Benoit Castel			
		NAME			
		35,041			
		REGISTRATION NO.			
BC/ia					
Date: April 1, 2002					

Application Data Sheet**Application Information**

Application Type:: Regular
Subject Matter:: Utility
Suggested Classification::
Suggested Group Art Unit::
CD-ROM or CD-R?: None
Number of CD disks::
Number of Copies of CDs::
Sequence Submission?: None
Computer Readable Form (CRF):: No
Number of copies of CRF:: 0
Title:: PLANT SEED ENDOSPERM-SPECIFIC
PROMOTERS
Attorney Docket Number:: 0512-1030
Request for Early Publication?: No
Request for Non-Publication?: No
Suggested Drawing Figure::
Total Drawing Sheets:: 11
Small Entity?: No
Latin Name::
Variety Denomination Name::
Petition Included?: No
Petition Type::
Licensed US Gov't Agency::
Contract or Grant Numbers::
Secrecy Order in Parent Appl.?: No

Applicant Information

Applicant Authority Type:: Inventor
 Primary Citizenship Country:: FRANCE
 Status:: Full Capacity
 Given Name:: JEAN-FRANCOIS
 Middle Name::
 Family Name:: BONELLO
 City of Residence:: BUSSY ST GEORGES
 State or Province of Residence::
 Country of Residence:: FRANCE
 Street of Mailing Address:: 9 RUE J.MONET

City of Mailing Address:: BUSSY ST GEORGES
 State or Province of Mailing Address::
 Country of Mailing Address:: FRANCE
 Postal or Zip Code of Mailing Address:: 77600

Applicant Authority Type:: Inventor
 Primary Citizenship Country:: GERMANY
 Status:: Full Capacity
 Given Name:: PETER
 Middle Name::
 Family Name:: ROGOWSKY
 City of Residence:: LYON
 State or Province of Residence::
 Country of Residence:: FRANCE
 Street of Mailing Address:: 31 RUE ANDRE BOLLIER

City of Mailing Address:: LYON
 State or Province of Mailing Address::
 Country of Mailing Address:: FRANCE
 Postal or Zip Code of Mailing Address:: 69007

Applicant Authority Type:: Inventor
 Primary Citizenship Country:: FRANCE
 Status:: Full Capacity
 Given Name:: PASCUAL
 Middle Name::
 Family Name:: PEREZ
 City of Residence:: CHANONAT
 State or Province of Residence::
 Country of Residence:: FRANCE
 Street of Mailing Address:: 17 CHEMIN DE LA PRADELLE
 VARENNES
 City of Mailing Address:: CHANONAT
 State or Province of Mailing Address::
 Country of Mailing Address:: FRANCE
 Postal or Zip Code of Mailing Address:: 63450

Correspondence Information

Correspondence Customer Number:: 000466

Representative Information

Representative Customer Number::	000466
----------------------------------	--------

Domestic Priority Information

Application::	Continuity Type::	Parent Application::	Parent Filing Date::
This application	National Stage of	PCT/FR00/02596	9/19/00

Foreign Priority Information

Country::	Application Number::	Filing Date::	Priority Claimed::
FRANCE	99 12305	10/1/99	Yes

Assignment Information

Assignee Name::

Street of Mailing Address::

City of Mailing Address::

State or Province of Mailing Address::

Country of Mailing Address::

Postal or Zip Code of Mailing Address::

10/089612

JC10 Rec'd PCT/PTO 01 APR 2002

PATENT
0512-1030

IN THE U.S. PATENT AND TRADEMARK OFFICE

In re application of: Jean-Francois BONELLO et al.

Appl. No.: NEW Group:

Filed: April 1, 2002 Examiner:

For: PLANT SEED ENDOSPERM-SPECIFIC PROMOTERS

PRELIMINARY AMENDMENTAssistant Commissioner for Patents
Washington, DC 20231

April 1, 2002

Sir:

The following preliminary amendments and remarks are respectfully submitted in connection with the above-identified application.

IN THE CLAIMS:

Please cancel claims 1-16 without prejudice or disclaimer of the subject matter contained therein.

Please add the following claims:

--17. (new) An isolated promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm

surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm.--

--18. (new) A promoter nucleotide sequence according to claim 17, isolated from cereals.--

--19. (new) A promoter nucleotide sequence according to claim 17, isolated from maize.--

--20. (new) A promoter nucleotide sequence according to claim 17, comprising a sequence chosen from amongst the sequences SEQ ID N° 1, N° 2, N° 3, N° 4, N° 5, N° 6 and N° 7, and any sequence which is a homologue of these.--

--21. (new) A promoter nucleotide sequence according to claim 17, also comprising a regulator element in *cis* defined by the pattern CTACACCA.--

--22. (new) A promoter nucleotide sequence according to claim 17, also comprising a regulator element in *cis* defined by the pattern CTACACCA, repeated in tandem.--

--23. (new) An expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of the endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--24. (new) An expression cassette according to claim 23, in which the gene of interest codes for a protein is chosen from amongst a protein involved in the biological characteristics selected from the group consisting of the development of the embryo, the development of the endosperm, the cell growth, the metabolism of sugars, the metabolism of the fatty acids, the metabolism of a toxic protein and the metabolism of a transcription inhibiting protein.--

--25. (new) An expression cassette according to claim 23, in which the gene of interest codes for a first protein is chosen from amongst barnase and isopentenyltransferase.--

--26. (new) An expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--27. (new) An angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular

in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--28. (new) An angiosperm plant host cell in the form of cereal transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--29. (new) A transgenic plant, selected from the group consisting of fruit, seed, grain and pollen, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--30. (new) A part of a transgenic plant, selected from the group consisting of fruit, seed, grain and pollen, generated from

an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--31. (new) A transgenic plant, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, wherein the transgenic plant is a plant selected from the group consisting of a cereal, an oily plant, maize, wheat, rape and sunflower.--

--32. (new) A part of a transgenic plant selected from the group consisting of fruit, seed, grain and pollen, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding

sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, wherein the transgenic plant is a plant selected from the group consisting of a cereal, an oily plant, maize, wheat, rape and sunflower.--

--33. (new) A hybrid transgenic plant obtained by crossing parts of transgenic plants, each part of the transgenic plant being selected from the group consisting of fruit, seed, grain and pollen, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--34. (new) A hybrid transgenic plant obtained by crossing transgenic plants, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is

bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, wherein the transgenic plant is a plant selected from the group selected from a cereal, an oily plant, maize, wheat, rape and sunflower.--

--35. (new) A method of obtaining an angiosperm plant having improved agronomic or nutritional qualities, comprising the steps consisting of:

- transforming at least one angiosperm plant cell by means of an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest; and

- cultivating the cell thus transformed so as to generate a plant containing in its genome an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression

being i) specific to the region of the endosperm surrounding the embryo in the seeds of the angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest.--

--36. (new) Use of an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, said use of the expression cassette being for obtaining a transgenic angiosperm plant exhibiting improved agronomic qualities.--

--37. (new) Use of an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, said use of the expression cassette being for obtaining a transgenic angiosperm plant exhibiting improved nutritional qualities.--

--38. (new) Use according to claim 36, for obtaining a transgenic plant producing seeds selected from the group consisting of starch contents modified compared with a non-transformed plant.--

--39. (new) Use according to claim 36, for obtaining a transgenic plant producing seeds selected from the group consisting of oil contents modified compared with a non-transformed plant.--

--40. (new) Use of a plant, selected from the group consisting of a seed, grain and fruit, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, the plant being for the preparation of derived products, notably food products.--

--41. (new) Use of a plant, selected from the group consisting of a seed, grain and fruit, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i)

specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, the part of the plant being for the preparation of derived products, notably food products.--

--42. (new) Use of a transgenic plant, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, wherein the transgenic plant is a plant selected from the group consisting of a cereal, an oily plant, maize, wheat, rape and sunflower, the plant being for the preparation of derived products, notably food products.--

--43. (new) Use of a part of a transgenic plant, selected from the group comprising a seed, grain and fruit, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i)

specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, the plant being for the preparation of derived products, notably food products.--

--44. (new) Use of a part of a hybrid transgenic plant, selected from the group comprising a seed, grain and fruit, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, the plant being for the preparation of derived products, notably food products.--

--45. (new) Use of a hybrid transgenic plant, generated from an angiosperm plant host cell, transformed by an expression vector containing an expression cassette comprising an isolator promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound, the said expression being i) specific to the region of an endosperm surrounding an embryo in seeds of angiosperms and ii) intervening in particular in the

Docket No. 0512-1030

early stages of development of the endosperm, the expression cassette being effectively bound to at least one gene of interest, wherein the hybrid transgenic plant is a plant selected from the group comprising a cereal, an oily plant, maize, wheat, rape and sunflower, the plant being for the preparation of derived products, notably food products.--

Docket No. 0512-1030

REMARKS

Claims 17-45 are pending in the present application. Claims 1-16 have been cancelled and claims 17-45 have been added.

Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly requested.

Should there be any matters that need to be resolved in the present application, the Examiner is respectfully requested to contact the undersigned at the telephone number listed below.

The Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 25-0120 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17.

Respectfully submitted,

YOUNG & THOMPSON



Benoit Castel, Reg. No. 35,041

745 South 23rd Street
Arlington, VA 22202
Telephone (703) 521-2297

BC/ia
Attachments

Atty. Docket No. 0512-1030

PATENTS

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Jean-Francois BONELLO et al.

Confirmation No. 2604

Serial No. 10/089,612

BOX PCT

Filed April 1, 2002

PLANT SEED ENDOSPERM-SPECIFIC
PROMOTERS

PRELIMINARY AMENDMENT

Commissioner for Patents

Washington, D.C. 20231

Sir:

Prior to the first Official Action, please amend the
above-identified application as follows:

IN THE SPECIFICATION:

Replace the paragraph beginning at page 2, line 29,
with the following rewritten paragraph:

--The promoter sequences of invention can
advantageously be selected from the group consisting of the
sequence comprising the sequences SEQ ID NO: 1, NO: 2, NO: 3, NO:
4, NO: 5, NO: 6 or NO: 7 and any nucleotide sequence which is a
homologue of these.--

Replace the paragraph beginning at page 3, line 1, with
the following rewritten paragraph:

--The sequence SEQ ID NO: 1 corresponds to the gene
promoter *Esrl*.--

BONELLO et al. S.N. 10/089,612

Replace the paragraph beginning at page 3, line 5, with the following rewritten paragraph:

--The sequence SEQ ID NO: 2 corresponds to the gene promoter *Esr2*.--

Replace the paragraph beginning at page 3, line 8, with the following rewritten paragraph:

--The sequence SEQ ID NO: 3 corresponds to the gene promoter *Esr3*.--

Replace the paragraph beginning at page 3, line 11, with the following rewritten paragraph:

--The sequence SEQ ID NO: 4 corresponds to the gene promoter *Esr4*.--

Replace the paragraph beginning at page 3, line 14, with the following rewritten paragraph:

--The sequence SEQ ID NO: 5 corresponds to a fragment of 499 pairs of bases on SEQ ID NO: 2 (nucleotides 1995-2493).--

Replace the paragraph beginning at page 3, line 17, with the following rewritten paragraph:

--The sequence SEQ ID NO: 6 corresponds to a fragment of 507 pairs of bases on SEQ ID NO: 3 (nucleotides 1202-1708).--

Replace the paragraph beginning at page 3, line 20, with the following rewritten paragraph:

--The sequence SEQ ID NO: 7 is a consensus sequence of 265 nucleotides, obtained by means of comparison between the sequences SEQ ID NO: 1, NO: 2 and NO: 3.--

BONELLO et al. S.N. 10/089,612

Replace the paragraph beginning at page 3, line 24, with the following rewritten paragraph:

--"Homologous nucleotide sequence" means any nucleotide sequence which differs from the sequence SEQ ID NO: 1, NO: 2, NO: 3, NO: 4, NO: 5, NO: 6 or NO: 7, by a substitution, deletion and/or insertion of one or more nucleotides, at positions such that these homologous nucleotide sequences preserve the property of specific promoter of the sequences SEQ ID NO: 1 to NO: 7.--

Replace the paragraph beginning at page 3, line 32, with the following rewritten paragraph:

--Preferably such a homologous nucleotide sequence is identical to at least 70% of the sequences SEQ ID NO: 1 to NO: 7, preferably at least 80%, preferably still at least 95%.--

Replace the paragraph beginning at page 4, line 18, with the following rewritten paragraph:

--Preferentially, such homologous nucleotide sequences specifically hybridises to the sequences which are complementary to the sequences SEQ ID NO: 1 to NO: 7 under stringent conditions. The parameters defining the stringency conditions depend on the temperature at which 50% of the paired strands separate (T_m).--

Replace the paragraph beginning at page 5, line 9, with the following rewritten paragraph:

--The various nucleotide sequences of the invention can be of artificial origin or not. They may be DNA sequences

BONELLO et al. S.N. 10/089,612

obtained by sieving banks of sequences by means of sensors produced on the basis of the SEQ ID NO: 1 to NO: 7. Such banks can be prepared by conventional techniques of molecular biology, known to persons skilled in the art.--

Replace the paragraph beginning at page 13, line 9, with the following rewritten paragraph:

--Figure 9 shows a comparison of the sequences of the promoters of the genes *Esr1*, *Esr2* and *Esr3* (respectively prEsr1, prEsr2 and prEsr3, or residues 271 to 531 of SEQ ID NO: 1, residues 1992 to 2493 of SEQ ID NO: 2 and residues 1199 to 1708 of SEQ ID NO: 3), the preserved parts being aligned.--

Replace the paragraph beginning at page 14, line 25, with the following rewritten paragraph:

--As illustrated in Figure 1, fragments containing the open reading phases of *Esr1*, *Esr2* and *Esr3* (Opsahl et al., 1997) as well as upstream and downstream sequences were sub-cloned in the plasmid pBluescript SK+ (Stratagene) in accordance with the methods described in Sambrook et al. (1989). The result was the plasmid L23/7 containing a fragment Sall of 2.1 kb of λ Esr1g1 and the plasmid L33/1 containing a fragment BamHI of 3.4 kb of λ Esr2g1. The plasmid L33/10 containing a fragment BamHI of 1.9 kb of λ Esr2g1 and the plasmid L102c24 containing a fragment HindIII of 4.5 kb of λ E1-111. XbaI sites situated just upstream of the open reading phase (TCTAGATTCCATG) (SEQ ID NO: 22) made it

BONELLO et al. S.N. 10/089,612

possible to differentiate the putative promoters of the respective open reading phases. In particular, the fragment *SaII/XbaI* of 0.53 kb of L23/7 was designated as a putative promoter of *Esr1*, the fragment *HindIII/BamHI* of 2.35 kb of L33/1 (upstream part of the promoter) and the fragment *BamHI/XbaI* of 0.14 kb of L33/10 (downstream part of the promoter), that of the putative promoter of *Esr2*, the fragment *HindIII/XbaI* of 1.71 kb, that of the putative promoter of *Esr3* and the fragment *XbaI/XbaI* of 1.62 kb comprising the putative promoter of *Esr4*. A functional promoter *Esr2* of 2.49 kb was reconstructed from the fragment *HindIII/BamHI* of 2.35 kb of L33/1 and from the fragment *BamHI/XbaI* of 0.14 kb of L33/10, in a base plasmid of the pBSSK+ type (Stratagene). The orientation of the arrows in Figure 1 represents the orientation 5'-3'.--

Replace the paragraph beginning at page 16, line 4, with the following rewritten paragraph:

--The consensus sequence (SEQ ID NO: 7) was obtained after alignment of the three promoter nucleotide sequences and using Sequencher 3.1 software from Genes Codes Corporation (Ann Arbor, MI 48106).--

Replace the paragraph beginning at page 17, line 8, with the following rewritten paragraph:

--A homology is also observed between the proximal regions which extend over approximately 500 pairs of bases between the promoter of *Esr2* and that of *Esr3*, as defined by the

BONELLO et al. S.N. 10/089,612

sequences SEQ ID NO: 5 and NO: 6.--

Replace the paragraph beginning at page 19, line 5, with the following rewritten paragraph:

--* the numbering of the promoter *Esr2* is based on the sequence p*Esr2* presented in an annexe (SEQ ID NO: 2) which goes from HindIII (AAGCTT or A = position 1) to XbaI (TCTAGA or A = position 2493).--

Replace the paragraph beginning at page 20, line 1, with the following rewritten paragraph:

--Secondly, fragments of the promoter were amplified using the initiator ESRX (5'GGGGTCTAGACTGTGAAGCTATTTTCCA3' (SEQ ID NO: 8)) containing the restriction site XbaI (underlined) and ESRH1 (5'GGGGAAGCTTTACATTCTTGCCATAACATA3' (SEQ ID NO: 9)), ESRH2 (5'GGGGAAGCTTTTCATCAATAATGCCTCATT3' (SEQ ID NO: 10)) or ESRH3 (5'GGGGAAGCTTTAATTCTTACTTCCTATCT3' (SEQ ID NO: 11)) containing the HindIII restriction site (underlined). The amplification products digested by XbaI and HindIII replaced the entire promoter *Esr2* upstream of the gene of the β -glucuronidase in the plasmid L124/19.--

Replace the paragraph beginning at page 20, line 13, with the following rewritten paragraph:

--The deleted promoters associated with the β -glucuronidase gene were then cloned in a plasmid containing the luciferase gene under the control of the promoter of the rice actin. The latter was obtained by cloning the fragment XhoI/NcoI

BONELLO et al. S.N. 10/089,612

of the plasmid pAct1-F4 (Mc Elroy D. et al., Mol Gen Genet., 231: 150-160, 1991) corresponding to the promoter and first intron of the rice actin, in a plasmid of the pGP214 type containing the luciferase gene and the terminator of nopaline synthase (Twell D. et al., Development 109, 705-713, 1990), digested by SalI/NcoI. An adaptor containing the restriction sites SalI and NotI, formed in nucleotides 5'GGCCAGTCGACAAAGCGGCCGCATGCA3' (SEQ ID NO: 12) and 5'TCAGCTGTTTCGCCGGCGT3' (SEQ ID NO: 13) was introduced into the plasmid obtained, digested by NotI and PstI (plasmid L210). The fragments SalI/NotI containing the deleted promoters associated with the β -glucuronidase gene were cloned in the plasmid L210 digested by SalI and NotI.--

Replace the paragraph beginning at page 23, line 8, with the following rewritten paragraph:

--¹⁾ adaptor JFB34:5'TCGACTGCAGCCCA 3' (SEQ ID NO: 14)

3'GACGTCGGGTTCGA 5' (SEQ ID NO: 15)

²⁾ adaptor JFB56: 5'CTAGACCCGAATTCGC 3' (SEQ ID NO: 16)

3'TGGGCTTAAGCGCCGG 5' (SEQ ID NO: 17)--

Replace the paragraph beginning at page 27, line 19, with the following rewritten paragraph:

--In a plasmid derived from pJIT30 containing the promoter 35S, a multiple cloning site and the terminating sequence of the cabbage mosaic virus (Guerineau F. et al., Plant Mol Biol, 15: 127-136, 1990), an adaptor containing a SpeI site and formed by the oligonucleotides (5'GATCCACTAGTCCCG (SEQ ID NO:

BONELLO et al. S.N. 10/089,612

18)) and (5'AATTCGGGACTAGTG (SEQ ID NO: 19)) was inserted between the sites BamHI and EcoRI. The fragment EcoRI/SpeI of the plasmid L42 a14 (Opsahl-Ferstad et coll., 1997) was inserted in the plasmid described previously. The construction thus obtained contained the gene *Esr2* in antisense orientation under the control of the promoter 35S (plasmid L79 b5).--

Replace the paragraph beginning at page 28, line 8, with the following rewritten paragraph:

--The promoter 35S was eliminated in the plasmid L79 b5 by restriction by SacI and HindIII, and replaced by an adaptor containing the restriction site HindIII and NotI and formed by the oligonucleotides (5'AAGCTTTTTCGGCCGC (SEQ ID NO: 20)) and (5'TCGAGCGGCCGCAAAAAGCTTAGCT (SEQ ID NO: 21)). The promoter *Esr2* in the form of a fragment HindIII/NotI of 2.44 kb was introduced into this adaptor. The construction thus obtained contains the gene *Esr2* in antisense orientation under the control of its own promoter (plasmid L129/46 (cf Figure 11)).--

IN THE CLAIMS:

Amend claim 20 as follows:

20. (amended) A promoter nucleotide sequence according to claim 17, comprising a sequence chosen from amongst the sequences SEQ ID NO: 1, NO: 2, NO: 3, NO: 4, NO: 5, NO: 6 and NO: 7, and any sequence which is a homologue of these.

BONELLO et al. S.N. 10/089,612

REMARKS

The above changes in the specification and claims amend the present application so that it is commensurate in scope with the present Sequence Listing.

Attached hereto is a marked-up version of the changes made to the specification and claims. The attached page is captioned "VERSION WITH MARKINGS TO SHOW CHANGES MADE."

Respectfully submitted,

YOUNG & THOMPSON

By Philip A. DuBois
Philip A. DuBois
Agent for Applicants
Registration No. 50,696
745 South 23rd Street
Arlington, VA 22202
Telephone: 521-2297

September 18, 2002

BONELLO et al. S.N. 10/089,612

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE SPECIFICATION:

Paragraph beginning at page 2, line 29, has been amended as follows:

The promoter sequences of invention can advantageously be selected from the group consisting of the sequence comprising the sequences SEQ ID [N° 1, N° 2, N° 3, N° 4, N° 5, N° 6 or N° 7] NO: 1, NO: 2, NO: 3, NO: 4, NO: 5, NO: 6 or NO: 7 and any nucleotide sequence which is a homologue of these.

Paragraph beginning at page 3, line 1, has been amended as follows:

The sequence SEQ ID [N°] NO: 1 corresponds to the gene promoter *Esr1*.

Paragraph beginning at page 3, line 5, has been amended as follows:

The sequence SEQ ID [N°] NO: 2 corresponds to the gene promoter *Esr2*.

Paragraph beginning at page 3, line 8, has been amended as follows:

The sequence SEQ ID [N°] NO: 3 corresponds to the gene promoter *Esr3*.

Paragraph beginning at page 3, line 11, has been amended as follows:

The sequence SEQ ID [N°] NO: 4 corresponds to the gene

BONELLO et al. S.N. 10/089,612

promoter *Esr4*.

Paragraph beginning at page 3, line 14, has been amended as follows:

The sequence SEQ ID [N°] NO: 5 corresponds to a fragment of 499 pairs of bases on SEQ ID [N°] NO: 2 (nucleotides 1995-2493).

Paragraph beginning at page 3, line 17, has been amended as follows:

The sequence SEQ ID [N°] NO: 6 corresponds to a fragment of 507 pairs of bases on SEQ ID [N°] NO: 3 (nucleotides 1202-1708).

Paragraph beginning at page 3, line 20, has been amended as follows:

The sequence SEQ ID [N°] NO: 7 is a consensus sequence of 265 nucleotides, obtained by means of comparison between the sequences SEQ ID [N° 1, N° 2 and N° 3] NO: 1, NO: 2 and NO: 3.

Paragraph beginning at page 3, line 24, has been amended as follows:

"Homologous nucleotide sequence" means any nucleotide sequence which differs from the sequence SEQ ID [N° 1, N° 2, N° 3, N° 4, N° 5, N° 6 or N° 7] NO: 1, NO: 2, NO: 3, NO: 4, NO: 5, NO: 6 or NO: 7, by a substitution, deletion and/or insertion of one or more nucleotides, at positions such that these homologous nucleotide sequences preserve the property of specific promoter of the sequences SEQ ID [N° 1 to N° 7] NO: 1 to NO: 7.

promoters of the genes *Esr1*, *Esr2* and *Esr3* (respectively prEsr1, prEsr2 and prEsr3, or [SEQ ID N° 1, SEQ ID N° 2, and SEQ ID N° 3] residues 271 to 531 of SEQ ID NO: 1, residues 1992 to 2493 of SEQ ID NO: 2 and residues 1199 to 1708 of SEQ ID NO: 3), the preserved parts being aligned.

Paragraph beginning at page 14, line 25, has been amended as follows:

As illustrated in Figure 1, fragments containing the open reading phases of *Esr1*, *Esr2* and *Esr3* (Opsahl et al., 1997) as well as upstream and downstream sequences were sub-cloned in the plasmid pBluescript SK+ (Stratagene) in accordance with the methods described in Sambrook et al. (1989). The result was the plasmid L23/7 containing a fragment Sall of 2.1 kb of λ Esr1g1 and the plasmid L33/1 containing a fragment BamHI of 3.4 kb of λ Esr2g1. The plasmid L33/10 containing a fragment BamHI of 1.9 kb of λ Esr2g1 and the plasmid L102c24 containing a fragment HindIII of 4.5 kb of λ E1-111. XbaI sites situated just upstream of the open reading phase (TCTAGATTCCATG) (SEQ ID NO: 22) made it possible to differentiate the putative promoters of the respective open reading phases. In particular, the fragment SaII/XbaI of 0.53 kb of L23/7 was designated as a putative promoter of *Esr1*, the fragment HindIII/BamHI of 2.35 kb of L33/1 (upstream part of the promoter) and the fragment BamHI/XbaI of 0.14 kb of L33/10 (downstream part of the promoter), that of the putative promoter of *Esr2*, the fragment HindIII/XbaI of 1.71 kb,

BONELLO et al. S.N. 10/089,612

that of the putative promoter of *Esr3* and the fragment *XbaI/XbaI* of 1.62 kb comprising the putative promoter of *Esr4*. A functional promoter *Esr2* of 2.49 kb was reconstructed from the fragment *HindIII/BamHI* of 2.35 kb of L33/1 and from the fragment *BamHI/XbaI* of 0.14 kb of L33/10, in a base plasmid of the pBSSK+ type (Stratagene). The orientation of the arrows in Figure 1 represents the orientation 5'-3'.

Paragraph beginning at page 16, line 4, has been amended as follows:

The consensus sequence (SEQ ID [N°] NO: 7) was obtained after alignment of the three promoter nucleotide sequences and using Sequencher 3.1 software from Genes Codes Corporation (Ann Arbor, MI 48106).

Paragraph beginning at page 17, line 8, has been amended as follows:

A homology is also observed between the proximal regions which extend over approximately 500 pairs of bases between the promoter of *Esr2* and that of *Esr3*, as defined by the sequences SEQ ID [N° 5 and N° 6] NO: 5 and NO: 6.

Paragraph beginning at page 19, line 5, has been amended as follows:

* the numbering of the promoter *Esr2* is based on the sequence p*Esr2* presented in an annexe (SEQ ID [N°] NO: 2) which goes from *HindIII* (AAGCTT or A = position 1) to *XbaI* (TCTAGA or A = position 2493).

Paragraph beginning at page 20, line 1, has been amended as follows:

Secondly, fragments of the promoter were amplified using the initiator ESRX (5'GGGGTCTAGACTGTGAAGCTATTTTCCA3' (SEQ ID [N°] NO: 8)) containing the restriction site XbaI (underlined) and ESRH1 (5'GGGGAAGCTTTACATTCTTGCCATAACATA3' (SEQ ID [N°] NO: 9)), ESRH2 (5'GGGGAAGCTTTTCATCAATAATGCCTCATT3' (SEQ ID [N°] NO: 10)) or ESRH3 (5'GGGGAAGCTTTAATTCTTACTTCCTATCT3' (SEQ ID [N°] NO: 11)) containing the HindIII restriction site (underlined). The amplification products digested by XbaI and HindIII replaced the entire promoter *Esr2* upstream of the gene of the β -glucuronidase in the plasmid L124/19.

Paragraph beginning at page 20, line 13, has been amended as follows:

The deleted promoters associated with the β -glucuronidase gene were then cloned in a plasmid containing the luciferase gene under the control of the promoter of the rice actin. The latter was obtained by cloning the fragment XhoI/NcoI of the plasmid pAct1-F4 (Mc Elroy D. et al., Mol Gen Genet., 231: 150-160, 1991) corresponding to the promoter and first intron of the rice actin, in a plasmid of the pGP214 type containing the luciferase gene and the terminator of nopaline synthase (Twell D. et al., Development 109, 705-713, 1990), digested by SalI/NcoI. An adaptor containing the restriction sites SalI and NotI, formed in nucleotides 5'GGCCAGTCGACAAAGCGCCGCATGCA3' (SEQ ID [N°] NO:

12) and 5'TCAGCTGTTTCGCCGGCGT3' (SEQ ID [N°] NO: 13) was introduced into the plasmid obtained, digested by NotI and PstI (plasmid L210). The fragments SalI/NotI containing the deleted promoters associated with the β -glucuronidase gene were cloned in the plasmid L210 digested by SalI and NotI.

Paragraph beginning at page 23, line 8, has been amended as follows:

¹⁾ adaptor JFB34:5'TCGACTGCAGCCCA 3' (SEQ ID [N°] NO: 14)

3'GACGTCGGGTTCGA 5' (SEQ ID [N°] NO: 15)

²⁾ adaptor JFB56: 5'CTAGACCCGAATTCGC 3' (SEQ ID [N°] NO: 16)

3'TGGGCTTAAGCGCCGG 5' (SEQ ID [N°] NO: 17)

Paragraph beginning at page 27, line 19, has been amended as follows:

In a plasmid derived from pJIT30 containing the promoter 35S, a multiple cloning site and the terminating sequence of the cabbage mosaic virus (Guerineau F. et al., Plant Mol Biol, 15: 127-136, 1990), an adaptor containing a SpeI site and formed by the oligonucleotides (5'GATCCACTAGTCCCG (SEQ ID [N°] NO: 18)) and (5'AATTCGGGACTAGTG (SEQ ID [N°] NO: 19)) was inserted between the sites BamHI and EcoRI. The fragment EcoRI/SpeI of the plasmid L42 a14 (Opsahl-Ferstad et coll., 1997) was inserted in the plasmid described previously. The construction thus obtained contained the gene *Esr2* in antisense orientation under the control of the promoter 35S (plasmid L79 b5).

Paragraph beginning at page 28, line 8, has been amended as follows:

The promoter 35S was eliminated in the plasmid L79 b5 by restriction by SacI and HindIII, and replaced by an adaptor containing the restriction site HindIII and NotI and formed by the oligonucleotides (5'AAGCTTTTTCGCGCCGC (SEQ ID [N°] NO: 20)) and (5'TCGAGCGCCGCAAAAAGCTTAGCT (SEQ ID [N°] NO: 21)). The promoter *Esr2* in the form of a fragment HindIII/NotI of 2.44 kb was introduced into this adaptor. The construction thus obtained contains the gene *Esr2* in antisense orientation under the control of its own promoter (plasmid L129/46 (cf Figure 11)).

IN THE CLAIMS:

Claim 20 has been amended as follows:

20. (amended) A promoter nucleotide sequence according to claim 17, comprising a sequence chosen from amongst the sequences SEQ ID [N° 1, N° 2, N° 3, N° 4, N° 5, N° 6 and N° 7] NO: 1, NO: 2, NO: 3, NO: 4, NO: 5, NO: 6 and NO: 7, and any sequence which is a homologue of these.

SEQUENCE LISTING

<110> BONELLO, JEAN-FRANCOIS
ROGOWSKY, PETER
PEREZ, PASCUAL

<120> PLANT SEED ENDOSPERM-SPECIFIC PROMOTERS

<130> 0512-1030

<140> 10/089,612

<141> 2002-04-01

<150> FR 99 12305

<151> 1999-10-01

<150> PCT/FR00/02596

<151> 2000-09-19

<160> 22

<170> PatentIn Ver. 2.1

<210> 1

<211> 531

<212> DNA

<213> Zea mays

<400> 1

gatcattaag	gactaagcag	tctttttccc	tttcgggcttg	catcatcttt	agtcttcac	60
actattataa	gccgaagcta	ttaccccttg	gctatagctt	cgggtgtcat	ctttattatc	120
ttcggactat	gtcttcacct	tgtatacctt	tgtcttgggg	gaaaaccttc	atcctgaagc	180
cgaagctccc	tgtaataaatt	catatcatgc	taaaaataaa	tggtcagtc	tgtttttgag	240
gaccttcgga	agaggaaggc	cccccaacaa	gacgattaac	tagtattgtc	tcactgcatt	300
gttttttttg	cacttcacat	ataatgcctc	aatagcatac	ttcatttttag	gaactttatt	360
aaaactgtct	taaaataggg	ccaagtcata	aattcattca	aagtgactct	tcattttctta	420
cttccatctt	ttgggtggtt	tgtatatata	tatgttcatt	gttgagtgat	gttccctacac	480
cactacacca	cacgttagat	atatatacac	aaaatagctt	cactatctag	a	531

<210> 2

<211> 2493

<212> DNA

<213> Zea mays

<400> 2

aagctttttc	ggtgatgaag	cacctgtaat	acttaacagc	atgctgaaaa	caaatagtta	60
gctgtgtttt	tgaggacctt	cgggaagatga	aggcccccaa	cacatcccat	gcatcaagtc	120
cccatgactt	gcaaaaaagc	aaattttatc	aaaattttct	ataaaaacact	tgaaaacatt	180
tctctttttg	aaaagtgtag	agcactagca	actgtctact	aaaaagggtc	ccaaattttc	240
gggtataaca	atcgcattgt	aaataacaca	aagggaatcc	tactaagagc	agtaatttgg	300
ctaaaaaca	agtgcatt	ttaatgta	agggaatagg	agcatgcaat	acttgtgttc	360
tttcagggtt	ttgatgtcct	caaaaagtgtg	ccccctggg	gcagttgcaa	cactcaaaat	420
ctactcgtat	acataaagaa	acatgggcac	aaaataagaa	acaatactca	aattatgaaa	480
aagggttcaaa	tggtccctata	attattgtag	acatttttaga	atttatttta	gaccaaaacc	540
atttaaaattt	ggttttaaaat	gagtttagata	ttaatatatta	ttcagtttat	agttatttgg	600
gacattttatt	tacttaacta	taacttctag	ggttttaaaa	gtaaaattttg	ggtccctagt	660
tggaaactagc	tcagattgct	gggtgatttc	cataaaaagtc	gaggttcctt	tagcaaaaat	720

ccacggtgaa	caagggggag	ataggtgttg	accgatatct	ctaaatthttg	atcgttggac	780
ggcacatgga	tgtctcagat	taaatggttg	atgtgcaagc	gacgcgcaca	cgatggagga	840
atggcttcac	gacggtgggc	tactagagct	ggctacgtca	accaatggag	ggctcgggtca	900
aggtcaaaat	ttgttgccaa	gccactgttg	ctcacgatga	gtcgattgag	cacatatcaa	960
ggtcgagggg	caaccagagg	ggcaagatcg	atgggtgcagt	gggtttctcg	atggaagggg	1020
aaacttcggg	gagcaattca	agatttccta	tcattgtgacc	gggtcaggga	atgggcgcac	1080
gggttttggg	accttctggg	gcacatcatg	ttgctgtatc	gatgtcaagg	gagcattagg	1140
gttcacgagt	cagcgatgac	gggcatgggt	ggacttgtgt	caccatgggt	cgatcaacta	1200
gggacgatag	agctctatga	agtttcacaa	cttcctcaca	ctctagggat	catggtgaca	1260
aaggtgggga	ggacggggcg	tctctagtga	gggtggaatg	cagttctgtc	acgtgggaat	1320
agtggcggca	tcgcttgtaa	tgaataaaaag	gtgcttgggt	ggctgggaag	tgcaatatga	1380
gggaagtagt	tgggtgcgggg	atgttccttt	tataaggag	caccattgat	taatggaaga	1440
caatgacaca	aagggtgggt	cgacagttta	aagctcgaat	gctgctaggg	gtgctcaagg	1500
ttaaaagatc	aggcatcagg	gaggaaaggc	agggataaaa	tttctttact	ccagttgtgg	1560
ggtgatgggg	acaagggttag	tgctcaagca	agggagggcg	agttcagcgc	agagatgcct	1620
gttgtgacac	atgggggggg	gggaattgga	ggttgggggt	gaccagggtga	cgttatggcg	1680
tgaccagag	aagagaccca	ctgatgggga	aaaaagggtgc	caacagggtg	ggaccaagggt	1740
gtcagtgaact	caccgtgaca	tgttattgga	aagttacgtc	cggaatgggt	tgggcctgag	1800
tgatctaggc	tggctcgggc	actgtgctga	tcctttaatt	tctccattcc	caatttaagt	1860
tgaattttta	attcaaatca	aatgactcca	aatctctcca	aaattaccaa	aatatagaat	1920
atttagatga	atatgttggg	ggagtgtggg	ctccgctttt	ggtagtatg	tttgtataaa	1980
aataattttct	ctccttttgt	cacttccaat	attgacttaa	atttttatgt	agcaatgcca	2040
acttttttta	gtagtgtgcc	acttatagca	caaaaactat	atccattttc	taatagtcct	2100
tgaaatccac	attctatttt	tagccattct	tcaaaattgg	cacaaaacta	ggaaaattta	2160
atacattctt	gccataacat	attctagtgc	aaatgttaac	tagattgtct	aatattagca	2220
aacttctttt	gtaagattca	ttaatatgtc	tacattgcat	acttttttag	aagttcatca	2280
ataatgctc	attagcatac	ttcattttag	gaacttgatt	aaaaccgcct	taaaatagag	2340
ccaagtgcg	gatccattta	aagggtgattc	ttaatttctt	acttcctatc	tttgggtggc	2400
tatgtttata	tatgtgtggg	tgggtgaatg	atgttctctac	accactacac	cacacgttgg	2460
acatatatat	ggaaaatagc	ttcacagtct	aga			2493

<210> 3

<211> 1708

<212> DNA

<213> Zea mays

<400> 3

aagcttagaa	atttttaaaaa	aagccaggca	agcgttggtg	tgcaaagagc	taaaaattag	60
gaagacaaga	gaacacggca	agaaagcatg	ctaaatgtgc	tcgcggtgcg	ttcttattta	120
tacgtcfaat	acgttgcaag	tggtagggcc	ccacttgtca	ttgactattg	ctattctagc	180
aaaggggaag	tgtttttcgg	accttcggct	taaggccttc	gtccatatcg	caatctgaat	240
ttatcattct	aacaaattaa	tattgtgagg	ggctactggt	gggggccttc	gacttccgaa	300
ggctctcaaa	aactgggtta	acagtgtttc	tggagtataa	tgcataaaca	ggtatcttcg	360
ggtttgatc	agaactacaa	catgaagagg	cacaaagaac	acgaagggtg	gcgcagagcc	420
gaagctcacg	tgtaggagag	cttcggcacg	acagcagaaa	aagggaaccg	acttaaaagg	480
aaaggctatt	cagacctcga	tggatttcta	taggtcatta	gcaaatgtaa	agggcattgaa	540
tgtaatttta	catgggctgt	gtccttgcc	ataaatagat	gaacagtact	ctcgtactgt	600
tcacgctgac	ttggcattcg	ctttttgcat	cacgttgtga	cccttgcttt	ccttcaaac	660
gaaggtacat	ctataatttg	ttattgtgtt	attgtggata	tggtaatgca	aataaaaaata	720
agttgatgat	aatgtttata	ttatttttcg	tatttcata	atgaattcct	cctcatcatt	780
tattgtgctt	acgaagggtt	ttccttcaaa	atctttgtcc	ggaattcatt	atatccgaag	840
ggaaataatg	tctcgaagga	cgaaggactt	tgatatttaa	cacttttcat	gttgccttgt	900
tcttgactct	tagcatttga	gaacaagtcc	ccaacagctc	ctaagctctt	ctttgaagaa	960
acaactacta	gatgaagttt	ctccaaaagt	acgtccattg	aatggagtaa	agagtcattt	1020
gacctctcgg	aataaaaatta	aaatgagaat	aagtaagaat	aaaacacctc	tattatcaaa	1080
tctaggccat	acaaacattg	ggtattacta	aaaaatagct	aatgccatct	ttcaaacatt	1140
ggaagttaaa	accaaccaat	cctcactcat	tccaagaaa	tattggatca	tatttaacat	1200

3

tttgtgtcac	ttacaaaaat	ggcttaatct	tttatgcggc	aatgccaaac	tttttttagca	1260
gggtgccact	tgtaacatga	aaactataac	tatttttcaaa	tagtaccttg	aaattcgcac	1320
tctattttta	tgcatctctc	aaaattgaca	caaattaaac	taggagaatt	caatacatctc	1380
ttgccataac	atattctaata	gcaaataatta	agtagattgc	tcaacatcgg	tacacatctt	1440
ttggacgatt	aattagtatt	gtctcactac	attctttgtt	ttagcagttc	atcaataatg	1500
cctcaatagc	atacttcatt	ttaggaactt	tatgaaaatt	gtcttaaaat	agggccaagt	1560
cacaaatcca	cttcaaaggt	gactcttcac	ttcttacttc	ctatctttgc	ttgtttttgt	1620
atatatatgt	gtggatgggt	gagtgatgtt	cctacaccac	tacaccacac	cttagacaca	1680
tatatggaaa	atagcttcac	tgtctaga				1708

<210> 4
 <211> 1232
 <212> DNA
 <213> Zea mays

<400> 4						
tagttcatgc	aaaagtagtg	agtgtttata	cacctatgcc	aatgaataac	ctctaactac	60
cctcatgtac	tggcctgggtg	tttggtaaac	atgaagcaat	aatttcctct	tatcaaaata	120
cgggtctcaa	gctgcatgtg	aaaaataata	aatatTTTTT	tgaagtgaat	gaaaaatgat	180
aaatatgaaa	cagtaaactc	ttccgttgaa	aaagtacatc	tctattatta	acgataacct	240
atatatcaat	ctacaatgcg	ctcatctgca	tctcgatgca	tactttcatc	atTTTTatgaa	300
tgtactttta	tgataagaag	gattagaatg	ttcttgtttt	cctcttattc	ttaccttttt	360
caaaattatc	agtttccaat	gtctgaatat	gcaatgcatt	ataaacctta	gtcagcatat	420
atcaagtcca	tatatataatgc	tatatgttta	agaactgggtg	ctgagtatgt	ctactcaaca	480
tatttttttag	ctattggatc	gagcagttta	gtaaaggtaa	actacattta	tctatcttca	540
agttgtattt	tcccaccctt	aaattatgaa	agggagtaac	gtccactcc	aactggtgaa	600
agggaaacaaa	tttggctctcc	ggactgattt	cattgggtgtg	tctctatttt	ttaaaaaac	660
aaaaaaacat	atttgttctc	tgaaaattga	taattaatta	atcataaatt	agggaaaaaa	720
ctatatgaaa	ctagttatag	ttttcttcta	aaattattgt	ctgtctgttg	gtgctctagt	780
tatagagtta	taacatgaaa	actatagcca	ttttcaaaata	gtgccttgaa	attcattttt	840
gttttagccat	tcttcaaaat	tgccacccaaa	ccaggagaat	ttattaataa	attcacgccca	900
taacatatctc	tagtccaaat	gttaagtggga	ttgctcaata	tcattatact	tcttttggac	960
gattcatttag	tactgccttg	ttgcataactt	tgtttttagca	gttcatcaat	aatgcttcac	1020
tagcataatt	catgttagga	acttgattaa	aactgcctta	agaacatggc	aaagtgataa	1080
atccacttca	aaggcaatta	ttaatTTCTT	acttccctatc	tttgggtgggt	tttgtatata	1140
tgtgtgtggg	tggttgagtg	atgcacacat	tttctacac	cactacatca	caccttggac	1200
atatatgtga	aaaaatagct	tcgcagtcta	ga			1232

<210> 5
 <211> 499
 <212> DNA
 <213> Zea mays

<400> 5						
ttttgtcact	tccaatattg	acttaaaattt	ttatgtagca	atgccaaactt	tttttagtag	60
tgtgccactt	atagcacaaa	aactatatcc	atTTTtcta	agtccttgaa	atccacattc	120
tatttttagc	cattcttcaa	aattggcaca	aaactaggaa	aatttaatac	attcttgcca	180
taacatatctc	tagtgcaaat	gttaactaga	ttgctcaata	ttagcaaaact	tcttttgtaa	240
gattcattaa	tattgtctaca	ttgcataactt	ttttagaagt	tcatcaataa	tgccctcatta	300
gcatacttca	tttttaggaac	ttgattaaaa	ccgccttaaa	atagagccaa	gtgacggatc	360
catttaaagg	tgattcttaa	tttcttactt	cctatctttg	gtggcttatg	tttatatatg	420
tgtgggtggg	tgaatgatgt	tcctacacca	ctacaccaca	cgttggacat	atatatggaa	480
aatagcttca	cagtctaga					499

<210> 6
 <211> 507
 <212> DNA
 <213> Zea mays

<400> 6
 ttgtgtcact tacaaaaatg gcttaatctt ttatgcggca atgccaacct ttttttagcag 60
 ggtgccactt gtaacatgaa aactataact attttcaaag agtaccttga aattcgcat 120
 ctatttttat gcattcttca aaattgacac aaattaaact aggagaattc aatacattct 180
 tgccataaca tatttctaag caaatattaa gtagattgct caacatcggt acacatcttt 240
 tggacgatta attagtattg tctcactaca ttctttggtt tagcagttca tcaataatgc 300
 ctcaatagca tacttcattt taggaacttt atgaaaattg tcttaaaata gggccaagtc 360
 acaaatccac ttcaaagggtg actcttcatt tcttacttcc tatctttgct tgtttttgta 420
 tatatatgtg tggatgggtg agtgatgttc ctacaccact acaccacacc ttagacacat 480
 atatggaaaa tagcttcaat gtctaga 507

<210> 7
 <211> 265
 <212> DNA
 <213> Zea mays

<220>
 <221> modified_base
 <222> (1)
 <223> a, c, t, g, other or unknown

<220>
 <221> modified_base
 <222> (32)
 <223> a, c, t, g, other or unknown

<220>
 <221> modified_base
 <222> (35)
 <223> a, c, t, g, other or unknown

<220>
 <221> modified_base
 <222> (128)
 <223> a, c, t, g, other or unknown

<220>
 <221> modified_base
 <222> (136)
 <223> a, c, t, g, other or unknown

<220>
 <221> modified_base
 <222> (175)
 <223> a, c, t, g, other or unknown

<400> 7
 namgattmay tartattggy wcaatrcatw snttnttttr gmasttcatc aataatgcct 60
 cawtagcata cttcatctta ggaacttkat kaaaaygyt ttaaaatagr gccaagtsay 120
 rratycantt yaaagntgay tcttmatttc ttacttctta tctttgstkg yttwngtwt 180
 tataatrtgk srtgggtgar tgatgttctt acaccactac accacacstt rgayayatat 240
 aayrgaaaata gcttcacwrt ctaga 265

<210> 8
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 8
gggtctaga ctgtgaagct attttcca

28

<210> 9
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 9
ggggaagctt tacattcttg ccataacata

30

<210> 10
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 10
ggggaagctt ttcataata atgcctcatt

30

<210> 11
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 11
ggggaagctt taatttctta cttcctatct

30

<210> 12
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 12
 ggccagtcga caaagcggcc gcatgca 27

<210> 13
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 13
 tcagctgttt cgccggcgt 19

<210> 14
 <211> 14
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 14
 tcgactgcag ccca 14

<210> 15
 <211> 14
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 15
 agcttgggct gcag 14

<210> 16
 <211> 16
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 16
ctagacccga attcgc 16

<210> 17
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 17
ggccgcgaat tcgggt 16

<210> 18
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 18
gatccactag tcccg 15

<210> 19
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 19
aattcgggac tagtg 15

<210> 20
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 20
aagctttttg cggccgc 17

<210> 21
<211> 25

8

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 21
tcgagcggcc gcaaaaagct tagct

25

<210> 22
<211> 13
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 22
tctagattcc atg

13

PLANT SEED ENDOSPERM-SPECIFIC PROMOTERS

The present invention relates to controlling the expression of genes during the development of the endosperm. It concerns in particular promoter nucleotide sequences which enable expression which is both specific to the interface between the embryo and the endosperm and early during the development of the endosperm.

10

The endosperm, a characteristic formation of Angiosperm seeds, is a nutritive tissue for the embryo. This is a tissue which is complex in its structure and development, in particular with cereals. The central area of the endosperm consists of large cells with vacuoles, which store the reserves of starch and proteins, whilst the region surrounding the embryo is distinguished by rather small cells, occupied for the major part by cytoplasm. At the present time the function of these cells, referred to as "dense cytoplasmic cells" (Schel et al. 1984) is not known. In 1994 Opsahl et al. identified a gene expressed specifically in this small region around the maize embryo, a gene which they called *Esr* standing for "Embryo Surrounding Region".

25

The authors of the present invention have now isolated promoter nucleotide sequences enabling an expression of the coding sequences with which they can be bound, which is specific to the region of the endosperm surrounding the embryo in Angiosperm seeds and which intervenes particularly in the early stages of the development of the endosperm.

The sequence SEQ ID N° 1 corresponds to the gene promoter *Esr1*.

5 The sequence SEQ ID N° 2 corresponds to the gene promoter
Esr2.

The sequence SEQ ID N° 3 corresponds to the gene promoter *Esr3*.

10

The sequence SEQ ID N° 4 corresponds to the gene promoter *Esr4*.

The sequence SEQ ID N° 5 corresponds to a fragment of 499
15 pairs of bases on SEQ ID N° 2 (nucleotides 1995-2493).

The sequence SEQ ID N° 6 corresponds to a fragment of 507 pairs of bases on SEQ ID N° 3 (nucleotides 1202-1708).

20 The sequence SEQ ID N° 7 is a consensus sequence of 265
nucleotides, obtained by means of comparison between the
sequences SEQ ID N° 1, N° 2 and N° 3.

25 "Homologous nucleotide sequence" means any nucleotide
sequence which differs from the sequence SEQ ID N° 1, N° 2,
N° 3, N° 4, N° 5, N° 6 or N° 7, by a substitution, deletion
and/or insertion of one or more nucleotides, at positions
such that these homologous nucleotide sequences preserve
the property of specific promoter of the sequences SEQ ID
30 N° 1 to N° 7.

Preferably such a homologous nucleotide sequence is identical to at least 70% of the sequences SEQ ID N° 1 to

N° 7, preferably at least 80%, preferably still at least 95%.

Homology is generally determined using a sequence analysis software (for example, the Sequence Analysis Software package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). Similar nucleotide sequences are aligned in order to obtain the maximum degree of homology (i.e. identity). To this end, it may be necessary to artificially introduce gaps in the sequence. Once the optimum alignment has been achieved, the degree of homology (i.e. identity) is established by recording all the positions for which the nucleotides of the two compared sequences are identical, with respect to the total number of positions.

Preferentially, such homologous nucleotide sequences specifically hybridises to the sequences which are complementary to the sequences SEQ ID N° 1 to N° 7 under stringent conditions. The parameters defining the stringency conditions depend on the temperature at which 50% of the paired strands separate (T_m).

For sequences comprising more than 30 bases, T_m is defined by the equation: $T_m = 81.5 + 0.41 (\% G + C) + 16.6 \log (\text{concentration in cations}) - 0.63 (\% \text{formamide}) - (600/\text{number of bases})$ (Sambrook et al., Molecular Cloning, A Laboratory Manual, Coldspring Harbor Laboratory Press, 1989, pages 9.54-9.62).

For sequences with a length less than 30 bases, T_m is defined by the equation: $T_m = 4(G + C) + 2(A + T)$.

Under appropriate stringency conditions, to which the aspecific sequences do not hybridise, the hybridisation temperature is approximately 5° to 30°C, preferably 5° to 10°C below T_m, and the hybridisation buffers used are preferably solutions with a ionic strength such as a 6xSSC solution for example.

The various nucleotide sequences of the invention can be of artificial origin or not. They may be DNA sequences obtained by sieving banks of sequences by means of sensors produced on the basis of the SEQ ID N° 1 to N° 7. Such banks can be prepared by conventional techniques of molecular biology, known to persons skilled in the art.

The nucleotide sequences according to the invention can also be prepared by chemical synthesis, or by mixed methods including the chemical or enzymatic modification of sequences obtained by sieving banks.

The promoter nucleotide sequences of the invention are preferably sequences isolated from cereals, in particular maize.

The promoter nucleotide sequences according to the present invention can in particular be isolated by methods of reversed PCR or working on the genome (Devic et al., 1997).

The promoter nucleotide sequences of the invention can also comprise or be associated with a *cis* CTACACCA regulating pattern, preferably repeated in tandem, or any

The gene of interest can be placed in sense or antisense orientation.

The promoter nucleotide sequence of the invention can also
5 be associated with a marker gene, for example a gene making it possible to select a plant transformed from a plant which does not contain transfected foreign DNA. As a marker gene, it is possible to cite in particular a gene confirming resistance to an antibiotic (Herrera-Estrella
10 et al., EMBO J. 2, 987-995 (1983)) or resistance to a herbicide (EP 242 246).

Another object of the invention is any nucleotide vector, such as a plasmid, which can be used for transforming host
15 cells, characterised in that it comprises an expression cassette as defined above. The construction of expression vectors for the transformation is within the capability of one skilled in the art following standard techniques.

20 Another object of the invention is an Angiosperm plant host cell, notably a cereal, transformed by a vector according to the invention.

The invention also concerns a transgenic plant or part of
25 a transgenic plant, in particular seed, fruit or pollen, generated from such a cell.

Amongst the cells able to be transformed according to the method of the invention, examples are cells of extensively
30 farmed plants (maize, wheat, rape, sunflower, peas, soya, barley, etc.) or food plants and flowers. Preferentially, it is possible to choose plants known to contain large

reserves (protein, glucidic and lipidic), in particular cereal plants and oily plants.

5 The hybrid plants obtained by crossing plants according to the invention also form part of the invention.

Another object of the invention is a method of obtaining an Angiosperm plant having improved agronomic or nutritional qualities, comprising the steps consisting of:

10

- transforming at least one Angiosperm plant cell by means of a vector as defined previously;

15

- cultivating the cell thus transformed so as to generate a plant containing in its genome an expression cassette according to the invention.

The transformation of vegetable cells can be achieved by the techniques known to one skilled in the art.

20

It is possible to cite in particular the methods of direct transfer of genes such as direct micro-injection into plant embryoids (Neuhaus et coll. 1997), vacuum infiltration (Bechtold et al. 1993) or electroporation (Chupeau et coll., 1989) or direct precipitation by means of PEG (Schocher et coll., 1986) or the bombardment by gun of particles covered with the plasmidic DNA of interest (Fromm M et al., 1990).

30

It is also possible to infect the plant with a bacterial strain, in particular Agrobacterium. According to one embodiment of the method of the invention, the vegetable cells are transformed by a vector according to the

invention, the said cell host being able to infect the said vegetable cells by allowing the integration, in the genome of the latter, of the nucleotide sequences of interest initially contained in the above-mentioned vector genome. Advantageously, the above-mentioned cell host used is *Agrobacterium tumefaciens*, in particular according to the method described in the article by An et al., (1986), or *Agrobacterium rhizogene*, in particular according to the method described in the article by Guerche et al. (1987).

For example, the transformation of vegetable cells can be achieved by the transfer of the T region of the tumour-inducing extra-chromosome circular plasmid of *Agrobacterium tumefaciens*, using a binary system (Watson et al., 1994). To do this, two vectors are constructed. In one of these vectors the T region has been eliminated by deletion, with exception of the right and left borders, a marker gene being inserted between them to allow selection in the plant cells. The other partner of the binary system is an auxiliary plasmid Ti, a modified plasmid which no longer has any T region but still contains the virulence genes *vir* necessary to the transformation of the vegetable cell.

25

According to a preferred mode, it is possible to use the method described by Ishida et al. (1996) for the transformation of Monocotyledons.

30 According to another protocol, the transformation is achieved according to the method described by Finer et al. (1992) using the tungsten or gold particle gun.

Another object in the invention is the use of the promoter nucleotide sequences referred to previously in molecular constructions intended to improve the agronomic, food or industrial quality of a plant, by acting in particular on
5 the size of the embryo or of the endosperm and/or its development.

This is because an early specific action on the development of the tissues of the embryo and of the
10 endosperm can be sought: according to the relative size of one or other tissue, it would be possible to obtain seeds or fruits with a higher starch (large endosperm) and/or oil (large embryo) content, via the use respectively of stimulator genes (hormone of the cellular
15 cycle for example) or inhibitor genes (toxic protein or transcription inhibitor for example). Endosperms without embryos could also be obtained according to this model, for industrial applications in starch making and semolina processing.

20

By way of example, the use of genes coding for hormones (cytokinins, auxins) of the cell cycle, under the control of the promoters described according to the invention, would make it possible to modify the processes of
25 cellularisation and, in a correlated fashion, the development of the endosperm in the light of the work of R J Scott (1998).

Action on the accumulation of nutrients in the embryo and
30 endosperm can also be sought, using for example, as genes of interest, genes coding for transporters of nutrients (sugar in particular), to the interfaces between mother plant/endosperm and endosperm/embryo, or genes coding for

inhibitors of these transports, for a differential accumulation of nutrients in the endosperm or embryo.

5 The invention therefore also relates to methods for
modifying the agronomic and/nutritional qualities of a
plant, through an early targeted action on the development
of the embryo/endosperm, using the transformation of the
plants with a vector according to the invention. In
particular, it is concerned with the modification of the
10 size and/or the development of the embryo/endosperm. It
also relates to the alteration of the development of the
embryo, with a view to producing seeds without embryos for
cereals in particular, presenting an interest for the
starch and semolina industries.

15

The object of the invention is more precisely the use of
an expression cassette as defined previously, for
obtaining a transgenic Angiosperm plant exhibiting
improved agronomic or nutritional qualities.

20

Advantageously, the transgenic plant obtained can produce
grains with starch or oil contents which are modified in
comparison with a non-transformed plant.

25 The invention also concerns the use of the transgenic
plants obtained according to the invention, or parts of
these plants in particular seeds, grains and fruits for
preparing derived products, in particular food products.

30 The products obtained, whether it be seeds with a higher
oil content, flours of seeds or grains with a higher
starch or oil content, also come within the scope of the
invention.

Finally, the object of the invention is any composition for human or animal food prepared from the said products obtained.

5

The following figures and examples illustrate the invention without limiting its scope.

LEGEND TO THE FIGURES

10

- Figure 1 depicts a diagram illustrating the steps of the cloning of the *Esr* promoters.

15

- Figure 2 depicts the restriction map of the plasmid L82/34, comprising in particular the promoter *pEsr1* fused with *Gus*.

20

- Figure 3 depicts the restriction map of the plasmid L124.19, comprising in particular the promoter *pEsr2* fused with *Gus*.

25

- Figure 4 depicts the restriction map of the plasmid L127a5, comprising in particular the promoter *pEsr3* fused with *Gus*.

- Figure 5 depicts the restriction map of the plasmid L78a1, comprising in particular the promoter *pEsr1* fused with *lpt*.

30

- Figure 6 depicts the restriction map of the plasmid L125a2, comprising in particular the promoter *pEsr2* fused with *lpt*.

- Figure 7 depicts the restriction map of the plasmid L77a101, comprising in particular the promoter *pEsr1* fused with *Barnase*.
- 5 - Figure 8 depicts the restriction map of the plasmid L126a3, comprising in particular the promoter *pEsr1* fused with *Barnase*.
- Figure 9 shows a comparison of the sequences of the
10 promoters of the genes *Esr1*, *Esr2* and *Esr3* (respectively *prEsr1*, *prEsr2* and *prEsr3*, or SEQ ID N° 1, SEQ ID N° 2, and SEQ ID N° 3), the preserved parts being aligned.
- 15 - Figure 10 depicts the restriction map of the plasmid pWP280.
- Figure 11 depicts the restriction maps of the plasmid L129/46, comprising notably the promoter *pEsr2* fused
20 with the sequence *Esr2* in antisense orientation.

EXAMPLES

EXAMPLE 1:

25

Quantitative expression *Esr1*, 2, 3.

The work by Opsahl-Ferstad et al. (1997) identified by "differential display" a specific amplicon of the
30 endosperm *Esr1* (access number on the EMBL database: X98495) and isolated by screening complementary genome banks on hybrid line HD5*HD7 (Barloy et coll. 1989) and line A188 (Gerdes and Tracey, 1993) respectively, of the

corresponding clones. From genome sequences *Esr1g1* (access number on EMBL: X98497) and *Esr2g1* (access number on EMBL: X98499) and *Esr3g2* (access number on EMBL: X99970) in particular, 3 genes *Esr1*, *Esr2* and *Esr3* were
5 revealed.

The authors of the present invention assessed the relative contributions of expression of each of the genes *Esr* by means of RT-PCR experiments, digestion by restriction
10 enzymes and quantification according to the methods known to persons skilled in the art, using DAP 7 and DAP 9 (day after pollination) equipment.

Quantification of the different bands identified on migration gel reveals relative contributions of 18%, 53% and 29% on average, for the transcripts of *Esr1*, *Esr2* and *Esr3* respectively. The promoter of *Esr2* therefore affords the strongest quantitative expression of the gene which it controls.
20

EXAMPLE 2:

Isolation and cloning of the promoter sequences

25 As illustrated in Figure 1, fragments containing the open reading phases of *Esr1*, *Esr2* and *Esr3* (Opsahl et al., 1997) as well as upstream and downstream sequences were sub-cloned in the plasmid pBluescript SK+ (Stratagene) in accordance with the methods described in Sambrook et al.
30 (1989). The result was the plasmid L23/7 containing a fragment *Sall* of 2.1 kb of λ *Esr1g1* and the plasmid L33/1 containing a fragment *BamHl* of 3.4 kb of λ *Esr2g1*. The plasmid L33/10 containing a fragment *BamHl* of 1.9 kb of

λEsr2g1 and the plasmid L102c24 containing a fragment HindIII of 4.5 kb of λE1-111. XbaI sites situated just upstream of the open reading phase (TCTAGATTCCATG) made it possible to differentiate the putative promoters of the
5 respective open reading phases. In particular, the fragment SalI/XbaI of 0.53 kb of L23/7 was designated as a putative promoter of Esr1, the fragment HindIII/BamHI of 2.35 kb of L33/1 (upstream part of the promoter) and the fragment BamHI/XbaI of 0.14 kb of L33/10 (downstream part
10 of the promoter) that of the putative promoter of Esr2, the fragment HindIII/XbaI of 1.71 kb, that of the putative promoter of Esr3 and the fragment XbaI/XbaI of 1.62 kb comprising the putative promoter of Esr4. A functional promoter Esr2 of 2.49 kb was reconstructed from the
15 fragment HindIII/BamHI of 2.35 kb of L33/1 and from the fragment BamHI/XbaI of 0.14 kb of L33/10, in a base plasmid of the pBSSK+ type (Stratagene). The orientation of the arrows in Figure 1 represents the orientation 5'-3'.

20

EXAMPLE 3:**Structure of the sequences upstream of the *Esr* genes**

25 Comparisons between the sequences of the regions 5' showed two types of homologies: a highly preserved sequence which corresponds to a proximal sequence of 265 pairs of bases and sequences of retrotransposons in the distal part. As the sequences of retrotransposons are in
30 different orientations and positions in the three promoters, they do not seem to fulfil a role in the expression of the *Esr* genes. Consequently, the 265 pairs of bases will contain all the *cis* information necessary

for an expression of specific genes of the region surrounding the embryo.

The consensus sequence (SEQ ID N° 7) was obtained after
5 alignment of the three promoter nucleotide sequences and using Sequencher 3.1 software from Genes Codes Corporation (Ann Arbor, MI 48106).

The degenerated bases are described in the Nomenclature
10 Committee of the International Union of Biochemistry (1985): Nomenclature for Incomplete Specified Bases in Nucleic Acid Sequences, European Journal of Biochemistry 150: 1-5.

15 In particular,

B = C, G or T but not A

D = A, G or T but not C

20

H = A, C or T but not G

K = G or T

25 M = A or C

N = G, A or C

R = G or A

30

S = C or G

V = A, C or G but not T

W = A or T

X = G, A, T or C

5

and Y = C or T.

A homology is also observed between the proximal regions which extend over approximately 500 pairs of bases between the promoter of *Esr2* and that of *Esr3*, as defined by the sequences SEQ ID N° 5 and N° 6.

The presence of elements acting in *cis* is sought amongst the preserved sequences, the most remarkable being CTACACCA, in tandem just 50 bases upstream of the open reading phase (Figure 9). This sequence is a good candidate for being an element responsible for a tissue-specific gene expression. The first of these repetitions is also placed in the loop of the greatest reversed repetition found in the three promoters. The sequences repeated more than twice are preserved only between the promoters *pEsr2* and *pEsr3*, in the missing region of *Esr1*: for example the sequences ATTCT and TTTA, each being repeated four times, a potential transcription enhancer in the light of the lowest expression of *Esr1* (Figure 9).

To demonstrate the functionality of the *cis* elements, constructs comprising deleted promoter nucleotide sequences, fused with *GUS*, were prepared.

30

By way of example, two techniques were used to create deletions of the promoter *Esr2*:

- by extensive digestion of 5' to 3' on the fragment HindIII-XbaI by means of the Erase-a-base™ kit from Promega (constructions L140);

- 5 - by PCR amplification of fragments of the promoter (constructions L194).

The plasmids L190 and L194 contain deleted promoter ESR2 fused with a reporter gene Gus and a terminator in accordance with the techniques described in the following example.

For the transformation, the fragments containing the constructs, deleted promoters constructs ESR2 - Gus - ter' were transferred into another plasmid containing the construct "promoter ubiquitin-luciferase-ter", the latter serving as an internal standard for quantifying the Gus activity and correcting the position effect of the insertion of the transgene in the genome on the expression, variable from one transformed plant to another.

The fragments of the promoter ESR2 resulting from these deletions are set out in the following table 1:

Table 1:

Chosen deletion technique	Remaining fragment of the promoter <i>Esr2</i> *
Digestion kit (L140)	985-2493
	1254-2493
	1865-2493
	1874-2493
	1880-2493
	2077-2493
Amplification PCR (L194)	2163-2493
	2275-2493
	2373-2493

5 * the numbering of the promoter *Esr2* is based on the sequence p*Esr2* presented in an annexe (SEQ ID N° 2) which goes from HindIII (AAGCTT or A = position 1) to XbaI (TCTAGA or A = position 2493).

10 To demonstrate the functionality of the promoter nucleotide sequences described above, the inventors cloned them upstream of the reporter gene GUS and used the constructs obtained for the transformation of plants.

15 In a preferred manner, the deleted promoters *Esr2* were obtained in accordance with the following protocols:

Firstly, deletions of 5' to 3' were effected using exonuclease III. The plasmid L124/19 containing the promoter of the gene *Esr2* coupled to the gene of the β -glucuronidase described in Example 4.1 was digested by HindIII in order to generate an initiation site for the deletions and by PstI to create a protection site against the action of the exonuclease III. The deletions were carried out with the Erase-a-base™ (Promega) kit.

Secondly, fragments of the promoter were amplified using the initiator ESRX (5'GGGGTCTAGACTGTGAAGCTATTTTCCA3' (SEQ ID N° 8)) containing the restriction site XbaI (underlined) and ESRH1 (5'GGGGAAGCTTTACATTCTTGCCATAACATA3' (SEQ ID N° 9)), ESRH2 (5'GGGGAAGCTTTTCATCAATAATGCCTCATT3' (SEQ ID N° 10)) or ESRH3 (5'GGGGAAGCTTTAATTTCTTACTTCCTATCT3' (SEQ ID N° 11)) containing the HindIII restriction site (underlined). The amplification products digested by XbaI and HindIII replaced the entire promoter *Esr2* upstream of the gene of the β -glucuronidase in the plasmid L124/19.

The deleted promoters associated with the β -glucuronidase gene were then cloned in a plasmid containing the luciferase gene under the control of the promoter of the rice actin. The latter was obtained by cloning the fragment XhoI/NcoI of the plasmid pAct1-F4 (Mc Elroy D. et al., Mol Gen Genet., 231: 150-160, 1991) corresponding to the promoter and first intron of the rice actin, in a plasmid of the pGP214 type containing the luciferase gene and the terminator of nopaline synthase (Twell D. et al., Development 109, 705-713, 1990), digested by SalI/NcoI. An adaptor containing the restriction sites SalI and NotI, formed in nucleotides 5'GGCCAGTCGACAAAGCGCCGCATGCA3' (SEQ ID N° 12) and 5'TCAGCTGTTTCGCCGGCGT3' (SEQ ID N° 13) was introduced into the plasmid obtained, digested by NotI and PstI (plasmid L210). The fragments SalI/NotI containing the deleted promoters associated with the β -glucuronidase gene were cloned in the plasmid L210 digested by SalI and NotI.

EXAMPLE 4:

Preparation of chimeric constructs

All the constructions can be effected in particular according to the methods described in Sambrook et al. (1989). The adaptors which can be used by way of example for cloning these fragments upstream of the different effecting genes are described in the restriction maps of the corresponding plasmids.

4-1 GUS chimeric constructs

The plasmid L23/7 (Esr1) was deleted from a fragment SacI containing undesirable restriction sites. Then a fragment XbaI/EcoRI of 2164 pairs of bases of the plasmid pBl101 (Jefferson et al., 1987) containing a *Gus* gene (coding for the β -glucuronidase but with no promoter) and a terminating sequence *nos*, was introduced. The new plasmid thus formed was then digested by XhoI and the digestion product containing the promoter region associated with the *Gus* gene and positioned upstream of the latter was subcloned in the vector pBCKS+ (Stratagene) so that the promoter is close the hybridisation zone of the initiator T7, thus enabling the plasmid L82/34 to be obtained (Figure 2, Table 2).

According to a similar protocol and with the restriction enzymes indicated in the corresponding figures, it was possible to obtain the plasmids L124/19 (pEsr2-GUS, Figure 3, Table 3) and L127a5 (pEsr3-GUS, Figure 4, Table 4).

It is also possible to use other reporter genes in replacement for GUS, for example GFP (Green Fluorescent

Protein, Siemering KR et al., 1996), to confirm the results obtained with GUS.

According to a protocol similar to that described previously, the fragment HindIII-XbaI of the promoter pEsr2 was fused with the coding sequence for GFP.

Table 2: Characteristics of the plasmid L82/34

Fragment	Position	Reference
pEsr1	741-1272	
Gus	1302-3107	
ter nos	3181-3434	
cat	5878-5223	
pBCKS+	1-740	Stratagene
L23/7 (pEsr1)	741-1272	Opsahl-Ferstad et al., 1997* and this example
pBI101	1273-3436	Jefferson et al., 1987
L23/7	3437-3763	Opsahl-Ferstad et al., 1997 and this example
(upstream)	3764-3769	Stratagene
pBSSK+	3770-6428	Stratagene
pBCKS+		

10

* the insert corresponds to the fragment Esr1g1 drawn in Figure 4 of Opsahl-Ferstad et al., 1997

Table 3: Characteristics of the plasmid L124/19

Fragment	Position	Reference
pEsr2	689-3175	
Gus	3205-5010	
ter nos	5084-5337	
bla	7441-6584	
pBSSK+	1-674	Stratagene
Linker JFB 34	675-688	this example ¹⁾
L33/1 (pEsr2')	689-3037	this example**
L33/10 (pEsr2'')	3038-3175	Opsahl-Ferstad et al., 1997 and this example**
pBI101	3176-5339	Jefferson et al., 1987
linker JFB56	5340-5346	this example ²⁾
pBSSK+	5347-7566	Stratagene

5 ** the inserts are presented partly as a fragment Esr2g1
in Figure 4 of Opsahl-Ferstad et al., 1997

1) adaptor JFB34: 5'TCGACTGCAGCCCA 3' (SEQ ID N° 14)

3'GACGTCGGGTTCGA 5' (SEQ ID N° 15)

10 2) adaptor JFB56: 5'CTAGACCCGAATTCGC 3' (SEQ ID N° 16)

3'TGGGCTTAAGCGCCGG 5' (SEQ ID N° 17)

Table 4: Characteristics of the plasmid L127a5

Fragment	Position	Reference
pEsr3	689-2390	
Gus	2420-4225	
ter nos	4229-4552	
bla	6656-5799	
Pbssk+	1-674	Stratagene
Linker JFB 34	675-688	this example
L102c24 (pEsr3)	689-2390	this example
pBI101	2391-4554	Jefferson et al., 1987
linker JFB56	4555-4561	this example
pBSSK+	4562-6781	Stratagene

5 4-2 Chimeric constructs lpt

The lpt gene codes for isopentenyl-transferase, which is an enzyme involved in the synthesis of cytokinin, a phytohormone implicated in vegetable cell growth. The gene sequence was determined by Heidekamp F. et al. (1983). Prior works also showed that this sequence, under the control of a specific promoter of the ovule, made it possible to increase the dry matter content in the fruit, in tomatoes, Martineau B. et al. (1995).

15

According to the cloning methods described above and with the fragments of nucleic acids and restriction enzymes indicated in the corresponding figures, it was possible to prepare constructs pEsr1-lpt (Figure 5, Table 5) and pEsr2-lpt (Figure 6, Table 6). It is also possible to obtain a construct pEsr3-lpt, according to the same protocol. For preparing these constructs, NcoI (CCATGG) sites straddling the codon ATG of the start of the open reading phase were used instead of the XbaI sites.

20

Table 5: Characteristics of the plasmid L78a1

Fragment	Position	Reference
pEsr1	310-844	
ipt	846-1565	
ter ipt	1566-	
bla	1850	
	2963-	
	3823	
pUC118	1-231	Boehringer
L23/7 (pEsr1)	232-844	Opsahl-Ferstad et al., 1997 and this example
isolated mutation	845	Zhang et al., 1996
pRZ1	846-1883	Zhang et al., 1995
pUC118	1884-	Boehringer
	4763	

5 Table 6: Characteristics of the plasmid L125a2

Fragment	Position	Reference
pEsr2	689-3183	
ipt	3185-3904	
ter ipt	3905-4189	
bla	6309-5449	
pBSSK+	1-674	Stratagene
Linker JFB 34	675-688	this example
L33/1 (pEsr2')	689-3037	this example
L33/10 (pEsr2'')	3038-3183	Opsahl-Ferstad et al., 1997 and this example
isolated mutation	3184	Zhang et al., 1996
pRZ1	3185-4198	Zhang et al., 1995
adaptor JFB56	4199-4214	this example
pBSSK+	4215-6434	Stratagene

4-3 Barnase chimeric constructs

- 10 The barnase gene codes for an Rnase. This gene was isolated using *Bacillus amyloliquefaciens* (Hartley, 1988). Its use for creating sterile male plants was described in

the application EP 344 029 published by Mariani et al. (1990).

In the context of the invention, the plasmids L77a101
 5 (pEsrl-barnase) and L126a3 (pEsr2-barnase) described in
 Figures 7 (Table 7) and 8 (Table 8) were obtained from the
 plasmid "promoter A6-barnase" described in WO 92/11379, by
 replacing pA6 with the promoters pEsrl and pEsr2
 respectively, in accordance with the techniques known to
 10 persons skilled in the art.

It is also possible to obtain a construct pEsr3-Barnase,
 in accordance with a similar protocol.

15 Table 7: Characteristics of the plasmid L77a101

Fragment	Position	Reference
pEsrl	80-605	
Barnase	613-945	
ter CaMV	1571-2257	
bla	4324-3467	
pA3	1-28	Scott et al., 1992
L23/7 (pEsrl)	29-605	Opsahl-Ferstad et al., 1997 and this example
pA3	606-4473	
		Scott et al., 1992

Table 8: Characteristics of the plasmid L126a3

Fragment	Position	Reference
pEsr2	43-2529	
Barnase	2537-2869	
ter CaMV	3495-4181	
bla	6248-5391	
pA3	1-28	Scott et al., 1992
linker JFB34	29-42	this example
L33/1 (pEsr2')	43-2391	this example
L33/10	2392-2529	Opsahl-Ferstad et al., 1997 and this example
(pEsr2'')	2530-6397	Scott et al., 1992
pA3		

5 4-4 Chimeric construct antiEsr2

The reconstituted functional promoter *Esr2* (2.49 kb), described in Example 2 and chosen preferentially in the light of the quantitative expression results described in Example 1, was fused with the sequence *Esr2g2* (Opsahl et al., 1997) taken in antisense orientation, itself fused with the terminator *Nos*.

In a preferred manner, the chimeric construct containing the gene *Esr2* in the reverse direction under the control of its own promoter was obtained in accordance with the following protocol:

In a plasmid derived from pJIT30 containing the promoter 35S, a multiple cloning site and the terminating sequence of the cabbage mosaic virus (Guerineau F. et al., Plant Mol Biol, 15: 127-136, 1990), an adaptor containing a *SpeI* site and formed by the oligonucleotides (5'GATCCACTAGTCCCG (SEQ ID N° 18)) and (5'AATTCGGGACTAGTG (SEQ ID N° 19)) was

inserted between the sites BamHI and EcoRI. The fragment EcoRI/SpeI of the plasmid L42 a14 (Opsahl-Ferstad et coll., 1997) was inserted in the plasmid described previously. The construction thus obtained contained the
5 gene *Esr2* in antisense orientation under the control of the promoter 35S (plasmid L79 b5).

The promoter 35S was eliminated in the plasmid L79 b5 by restriction by SacI and HindIII, and replaced by an
10 adaptor containing the restriction site HindIII and NotI and formed by the oligonucleotides (5'AAGCTTTTTCGGCCGC (SEQ ID N° 20)) and (5'TCGAGCGGCCGCAAAAAGCTTAGCT (SEQ ID N° 21)). The promoter *Esr2* in the form of a fragment HindIII/NotI of 2.44 kb was introduced into this adaptor.
15 The construction thus obtained contains the gene *Esr2* in antisense orientation under the control of its own promoter (plasmid L129/46 (cf Figure 11)).

According to a similar protocol, it is possible to obtain
20 the constructs comprising the promoter *Esr2* fused with the antisense sequences *Esr1* and *Esr3* respectively. It is also possible to obtain the same type of chimeric constructs with the other *Esr* promoters according to the invention.

25

Constructs comprising the constituent promoter 35S fused with the *Esr* antisense sequences described below have also been obtained.

30 **EXAMPLE 5:**

Obtaining transgenic plants (necessity for the stable transformation of maize)

Transient expression experiments using transformation by bombardment of vegetable cells, with chimeric constructs pEsr-GUS and constituent promoter-GUS respectively, did not give results revealing the specificity of expression of the promoters tested: no GUS activity was displayed in the area defined by the Esr cells. The small size of this area and other peculiar particularities could explain the fact that the technique is unsuited under standard conditions to transient expression. By way of example, the constituent promoters tested as a control are the rice actin promoters (McElroy et al., 1992), maize ubiquitin (Christensen et al., 1996), maize Adh (Dennis et al., 1984) and 35S (Odell et al., 1985), gave a blue colouring throughout the endosperm, demonstrating the functionality of the transformation system, but not in the area surrounding the embryo, which confirms the unsuitability of the system for this area.

The transformation aimed at a stable expression therefore became necessary for studying the specificity of expression of the promoters according to the invention.

5-1 Particle gun

The method used is based on the use of a particle gun identical to the one described by J. Finer (1992). The target cells are undifferentiated cells in rapid divisions which have preserved suitability for the regeneration of entire plants. This type of cell composes the embryogenic callus (referred to as type II) of maize. These calluses are obtained from immature embryos of the genotype Hill according to the method and on the media described by

Armstrong (Maize Handbook: 1994, M. Freeling, V. Walbot Eds, pp. 665-671). These fragments of the calluses with a surface area of 10 to 20 mm² were disposed, 4 hours before bombardment, at the rate of 16 fragments per dish, in the centre of a Petri dish containing a culture medium identical to the initiation medium, with 0.2 M of mannitol + 0.2 M of sorbitol added. The plasmids described in the previous examples and carrying the genes to be introduced are purified on a Qiagen^R column following the instructions of the manufacturer. They are then precipitated on particles of tungsten (M10) in accordance with the protocol described by Klein (1987). The particles thus coated are projected towards the target cells by means of the gun and in accordance with the protocol described by J. Finer (1992). The dishes of calluses thus bombarded are then sealed by means of Scellofrais^R and then cultivated in darkness at 27°C. The first planting out took place 24 hours afterwards, and then every fortnight for 3 months on a medium identical to the initiation medium with a selective agent added. After 3 months or sometimes earlier, calluses are obtained whose growth is not inhibited by the selective agent, normally and for the major part composed of cells resulting from the division of a cell which integrated in its genotype one or more copies of the selection gene. The frequency of obtaining such calluses is approximately 0.8 callus per dish bombarded.

These calluses are identified, individualised, amplified and then cultivated so as to regenerate plant germs, modifying the hormonal and osmotic balance of the cells in accordance with the method described by Vain et al. (1989). These plants are then acclimatised in a

greenhouse, where they can be crossed in order to obtain hybrids or self-fertilised.

5-2 Transformation by *Agrobacterium*

5

Another transformation technique which can be used in the context of the invention uses *Agrobacterium tumefaciens*, in accordance with the protocol described by Ishida et al. (1996), in particular from immature embryos from 10 days after fertilisation. All the media used are referenced in the reference cited. The transformation begins with a co-culture phase in which the immature embryos of the maize plants are brought into contact for at least 5 minutes with *Agrobacterium tumefaciens* LBA 4404 containing the superbinary vectors. The superbinary plasmid is the result of a homologous recombination between an intermediate vector carrying ADN-T containing the gene of interest and/or the selection marker derived from the plasmids described in the previous examples, and the vector pSB1 of Japan Tobacco (EP 672 752) which contains: the genes *virB* and *virG* of the plasmid pTiBo542 present in the supervirulent strain A281 of *Agrobacterium tumefaciens* (ATCC 37349) and a homologous region found in the intermediate vector allowing this homologous recombination. The embryos are then placed on a medium LSAs for 3 days in darkness and at 25°C. A first selection is effected on the transformed calluses. The embryogenic calluses are transferred onto a medium LSD5 containing phosphinotricine at 5 mg/l and cefotaxime at 250 mg/l (elimination or limitation of the contamination by *Agrobacterium tumefaciens*). This step is carried out 2 weeks in darkness and at 25°C. The second selection step is carried out by the transfer of the embryos which are

developed on an LSD5 medium, on an LSD10 medium (phosphinotricine at 10 mg/l) in the presence of cefotaxime, for 3 weeks under the same conditions as before. The third selection step consists of excising the
5 type I calluses (fragments of 1 to 2 mm) and transferring them 3 weeks in darkness and at 25°C onto an LSD 10 medium in the presence of cefotaxime.

The regeneration of the plant germs is carried out by
10 excising the type I calluses which have proliferated and transferring them onto an LSZ medium in the presence of phosphinotricine at 5 mg/l and cefotaxime for 2 weeks at 22°C and under continuous light.

15 The plant germs which have regenerated are transferred onto an RM + G2 medium containing 100 mg/l of Augmentin for 2 weeks at 22°C and under continuous illumination for the development step. The plants obtained are then transferred to the phytotron with a view to their
20 acclimatisation.

**5-3 Preferred mode for the barnase constructs:
retransformation of the act-barstar calluses**

25 The barnase chimeric constructs described in Example 3 can be used for conventional transformations according to one or other of the techniques described above.

According to a preferred mode, adapted to the toxic
30 character of barnase, pretransformed calluses are used for the transformation, containing the gene barstar, which codes for a specific inhibitor of Barnase (Hartley, 1988). This gene serves as "protection" during the process of

regenerating these calluses, which takes place essentially from embryogenesis in maize.

- step a: obtaining a line expressing barstar and a plasmid containing the gene for resistance to hygromycine:

A first transformation step is carried out in accordance with one of the protocols described, with the plasmid pWP280 containing the cassette pActin-intron-Barstar-Nos poly A.

This cassette was obtained according to the following steps: the barnase fragment was amplified with PCR from the plasmid pTG2 (Horovitz et al., 1990) and then subcloned as a fragment XbaI/HindIII in the plasmid pBluescript KS+ (Stratagene) giving the plasmid pWP118.

The barstar gene was then transferred as a fragment XbaI/HincII into a site XbaI/SmaI of the plasmid pW90, derived from the plasmid pJLT30 described by Guerineau et al. (1990) (promoter 35SCaMV replaced by the double promoter 35S and the *polylinker* region between the sites XbaI and EcoRI replaced by the sites SpeI, BamHI, SmaI and PstI).

The region polyA CaMV of the plasmid obtained is replaced by the region nos polyA of pED23 (Dale et al., 1991) forming the plasmid pWP266. Finally, the double promoter region 35S CaMV is replaced by the rice actin promoter and the intron derived from pCOR113 (Mc Elroy et al., 1991) forming the plasmid pWP280 (Figure 10).

The "actin-barstar promoter" plants thus produced are analysed by Northern Blot in order to identify the plants

correctly expressing ARNm coding for Barstar. The plants thus produced supply embryos expressing the Barstar gene, which will be used for producing type II calluses according to known techniques: putting the embryos in
5 culture on a medium inducing callogenesis and replanting on a selective medium containing hygromycin.

- step b: transformation of these calluses with the barnase chimeric construct:

10

The act-barstar calluses obtained at the previous step are then bombarded according to the technique described at point 5-1 with the "Esr-barnase promoter" construct previously described with a plasmid conferring resistance
15 to Basta (pDM302, Mc Elroy et al., 1991). The two genes are then separated into the descendants by segregation, in order to see the effect of the single promoter construct Esr-barnase.

20 Better results, particularly with regard to the effectiveness of transformation and the number of plants regenerated, were obtained according to this preferred mode, in comparison with the conventional technique which aims to transform the calluses directly by means of the
25 "Esr-barnase" constructs.

EXAMPLE 6

Demonstration of the functionality of the promoter
30 sequences (the case of GUS constructs)

In order to detect the β -glucuronidase activity, the maize seeds issuing from plants transformed by the particle gun

are harvested at precise stages of the development and cut along the longitudinal axis. They are incubated in the presence of 5-bromo-4-chloro-3-indolyl- β -D-glucuronic acid (X-GlcA, Duchefa), at 37°C for 24 hours (Jefferson et al.,
5 1987).

In the case of the construct Esr2-Gus in particular, the blue colouring is delimited at the contour of the embryo at the 4th and 5th days after pollination, and then only
10 at the suspensor level on the 6th and 7th days, and finally at the base of the suspensor at days 9, 12, 13 and 15.

The expression results in the transgenic plants demonstrate that the fragments 5' described in the present
15 invention correspond to functional promoters and that they are sufficient for a correct spatio-temporal expression, in accordance with the prior results of Opsahl et al. (1997).

20 The use of these promoter nucleotide sequences in molecular constructions intended to improve the agronomic, food or industrial quality of a plant is particularly advantageous for modifying the size of the embryo or of the endosperm and/or its development.

BIBLIOGRAPHY

- 5 - Armstrong (1994), *Maize Handbook*, Freeling, M., Walbot,
V. Eds, 665-671.
- Barloy, D. et coll. (1989), *Maydica*, 34, 303-308.
- 10 - Becker, H.A. et al. (1999), *Domains of gene expression*
in developing endosperm, 361-375.
- Breton, C. et al. (1995), *Plant Mol. Biol.*, 27, 105-113.
- 15 - Christensen et al. (1996), *Transgenic Res.*, 5: 213.
- Clark, J.K. and Sheridan, W.F. (1986), *J. Heredity*, 77,
83-92.
- 20 - Dale et al. (1990), *Gene*, 91: 79-85.
- Davis, R.W. et coll. (1990), *Can. J. Bot.*, 68, 471-479.
- 25 - Dennis (1884), *Nucl. Ac. Res.*, 12: 3983-4000.
- Devic et al. (1997), *Plant Physiol. Biochem.*, 35: 35(4):
331-339.
- Finer, J. (1992), *Plant Cell Report*, 11: 323-328.
- 30 - Gerdes, J.T. and Tracy, W.F. (1993), *Crop Sci.*, 33, 334-
337.
- Guerche et al. (1987), *Mol. Gen. Genet.*, 206: 382.

- Guerineau et al. (1990), *Plant. Mol. Biol.*, 15: 127-136.
- Hartley et al. (1988), *J. Mol. Biol.*, 202, 913-915.
- 5 - Heidekamp, F. et al. (1983), *Nucl Acids Res*, 11, 6211-6223.
- Horovitz et al. (1990), *J. Mol. Biol.*, 216: 1031-1044.
- 10 - Hu et al. (1995), *Molecular and General Genetics*, 248: 471-480.
- Hueros, G. et al. (1995), *Plant Cell*, 7, 747-757.
- 15 - Ishida et al. (1996), *Nature Biotechnology*, 14: 745-750.
- Jefferson et al. (1987), *Plant Molecular Biology Reporter*, 5(4): 387-405.
- 20 - Klein (1987), *Nature*, 327: 70-73.
- Kowles, R.V. and Phillips, R.L. (1988), *Int. Rev. Cytol.*, 112, 97-136.
- 25 - Kyle, D.J. and Styles, E.D. (1977), *Planta*, 137, 185-193.
- Liang, P. and Pardee, A.B. (1992), *Science*, 257, 967-971.
- 30 - Lopes, M.A. and Larkins, B.A. (1993), *Plant Cell*, 5, 1383-1399.

- Mariani et al. (1990), *Nature*, 347, 737-741.
- Mc Elroy et al. (1991), *Mol. Gen. Genet.*, 231: 150-160.
- 5 - McElroy et al., *Plant Cell*, 2: 163-171, 1990.
- Odell et al. (1985), *Nature*, 313: 810-812.
- 10 - Opsahl-Ferstad, H.D. et al. (1997), *The Plant J.*, 12, 235-246.
- Poole et al. (1985), *Cell*, 40: 37-43.
- 15 - Sambrook et al. (1989), *Molecular Cloning - A laboratory manual*, Cold Spring Harbor Laboratory Press.
- Schel, J.H.N. et coll. (1984), *Can. J. Bot.*, 62, 2842-2856.
- 20 - Scott et al. (1992), patent application WO 92/11 379.
- Siemering, K.R. et al. (1996), *Current Biology*, 6, 1653-1663.
- 25 - Twell, D. et al., *Development*, 109, 705-713, 1990.
- Vain et al. (1989), *Plant Cell Tissue and Organ Culture*, 18: 143-151.
- 30 - Xu, J. et al. (1995), *Plant Physiol.*, 108, 1293-1294.

- Zhang et al. (1995), The effect of auxin on cytokinin levels and metabolism in transgenic tobacco tissue an *ipt* gene, *Planta*, 196: 84-94.
- 5 - Zhang et al. (1996), Expression of the isopentenyl transferase gene is regulated by auxin in transgenic tobacco tissues, *Transgenic Res.*, 5: 57-65.

CLAIMS

1. An isolated promoter nucleotide sequence allowing an expression of the coding sequences to which it is bound;
5 said expression being i) specific to the region of the endosperm surrounding the embryo in the seeds of the angiosperms and ii) intervening in particular in the early stages of development of the endosperm.
- 10 2. The promoter nucleotide sequence according to Claim 1, isolated from cereals, in particular maize.
3. The promoter nucleotide sequence according to either one of Claims 1 or 2, comprising a sequence selected from
15 the group consisting of sequence SEQ ID N° 1, N° 2, N° 3, N° 4, N° 5, N° 6 or N° 7, and any sequence which is a homologue of these.
4. The promoter nucleotide sequence according to any one
20 of the preceding claims, also comprising a regulator element in *cis* defined by the pattern CTACACCA, preferably repeated in tandem.
5. An expression cassette comprising a promoter
25 nucleotide sequence according to any one of the preceding claims, operatively bound to at least one gene of interest.
6. Expression cassette according to Claim 5, in which the
30 gene of interest codes for a protein selected from the group consisting of a protein involved in the development of the embryo and/or of the endosperm, the cell growth,

the metabolism of the sugars or that of the fatty acids, a toxic protein, and a transcription inhibiting protein.

7. An expression cassette according to either one of
5 Claims 5 or 6, in which the gene of interest codes for a first protein chosen from amongst barnase or isopentenyltransferase.

8. An expression vector containing an expression cassette
10 according to any one of Claims 5 to 7.

9. An angiosperm plant host cell, particularly cereal, transformed by a vector according to Claim 8.

15 10. A transgenic plant or part of a transgenic plant, in particular fruit, seed, grain or pollen, generated from a cell according to Claim 9.

11. The plant or part of a plant according to Claim 10,
20 wherein said plant or part of plant is a cereal or oily plant, selected in particular from the group consisting of maize, wheat, rape and sunflower, preferentially maize.

12. A hybrid transgenic plant obtained by crossing plants
25 as defined in either one of Claims 10 or 11.

13. A method of obtaining an angiosperm plant having improved agronomic or nutritional qualities, comprising the steps consisting of:

30

- transforming at least one angiosperm plant cell by means of a vector according to Claim 8;

14. Use of an expression cassette as defined in any one of Claims 5 to 7, for obtaining a transgenic angiosperm plant exhibiting improved agronomic or nutritional qualities.

16. Use of a plant or part of a plant, in particular a
15 seed, grain or fruit, as defined in any one of Claims 10
to 12, for the preparation of derived products, notably
food products.

Fig. 1

Genome bank HD5 x HD7 (Esra sensor)

Fig. 9

Majority

5

10

Repetition CTACACC in tandem

Repetition TTTTA → Repetition ATTCT

Fig. 11

Plasmid

5

Promoter of Esr2 gene

Codified sequence of Esr2 gene

10 Terminator

LIST OF SEQUENCES

- 5 Specific promoters of the endosperm of vegetable seeds

Pages 4-7

Artificial sequence

5

Description of the artificial sequence: oligonucleotide

etc.

10

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION
EN MATIÈRE DE BREVETS (PCT)

(19) Organisation Mondiale de la Propriété
Intellectuelle
Bureau international



(43) Date de la publication internationale
12 avril 2001 (12.04.2001)

PCT

(10) Numéro de publication internationale
WO 01/25439 A1

- (51) Classification internationale des brevets⁷: C12N 15/29, 15/82, A01H 5/00
- (74) Mandataire: JACOBSON, Claude; Cabinet Lavoix, 2, place d'Estienne d'Orves, F-75441 Paris Cedex 09 (FR).
- (21) Numéro de la demande internationale:
PCT/FR00/02596
- (81) États désignés (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) Date de dépôt international:
19 septembre 2000 (19.09.2000)
- (25) Langue de dépôt: français
- (26) Langue de publication: français
- (30) Données relatives à la priorité:
99/12305 1 octobre 1999 (01.10.1999) FR
- (84) États désignés (*régional*): brevet ARIPO (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), brevet eurasien (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), brevet européen (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), brevet OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Déposant (*pour tous les États désignés sauf US*): BIO-GEMMA [FR/FR]; 1, rue Edouard-Colonne, F-75001 Paris (FR).
- (72) Inventeurs; et
- (73) Inventeurs/Déposants (*pour US seulement*): BONELLO, Jean-François [FR/FR]; 9, rue J. Monet, F-77600 Bussy St Georges (FR). ROGOWSKY, Peter [GE/FR]; 31, rue André Bollier, F-69007 Lyon (FR). PEREZ, Pascual [FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450 Chanonat (FR).
- Publiée:
— Avec rapport de recherche internationale.
- En ce qui concerne les codes à deux lettres et autres abréviations, se référer aux "Notes explicatives relatives aux codes et abréviations" figurant au début de chaque numéro ordinaire de la Gazette du PCT.

(54) Title: PLANT SEED ENDOSPERM-SPECIFIC PROMOTER

(54) Titre: PROMOTEURS SPECIFIQUES DE L'ALBUMEN DES GRAINES DE VEGETAUX

(57) Abstract: The invention concerns promoter nucleotide sequences enabling expression of encoding sequences whereto they can be bound, which is specific of the endosperm region enclosing the embryo of seeds of Angiosperms and which intervene in particular in the early stages of endosperm development, and their use for agronomic or nutritional improvement of plants.

(57) Abrégé: Cette invention concerne des séquences nucléotidiques promotrices permettant une expression des séquences codantes auxquelles elles peuvent être liées, qui est spécifique de la région de l'albumen entourant l'embryon dans les graines des Angiospermes et qui intervient en particulier dans les stades précoces du développement de l'albumen, ainsi que leur utilisation pour l'amélioration agronomique ou nutritionnelle des plantes.

WO 01/25439 A1

1/11




FIG.1




FIG.2

503	Pvu I
532	Pvu II
657	Kpn I
663	Apa I
688	Xho I
674	Sal I
683	Pst I
688	Hind III
730	Sph I
925	BatX I
1033	Sph I
1443	EcoR V
1675	Cla I
1779	Kpn I
1806	Cla I
1870	Nco I
1901	Sac I
303037	BamH I
31375	Xba I
3181	BamH I
3188	Sma I
3278	Nru I
3762	EcoR V
3800	Mlu I
3993	EcoR V
41495	Mlu I
4513	Mlu I
4919	Nru I
6072	Sac I
5086	Pvu I
5339	EcoR I
5348	Not I
5368	Sac II
5369	BatX I
5367	Sac I
5565	Pvu II
7024	Pvu I




FIG. 3

4/11




FIG.4

5/11




FIG.5

503 Pvu I
 532 Pvu I
 557 Kpn I
 583 Apa I
 608 Xho I
 634 Sal I
 660 Pst I
 686 Hind III
 712 Sph I
 738 BatX I
 764 Sph I
 790 Sph I
 816 EcoR V
 842 Cla I
 868 Cla I
 894 Kpn I
 920 Cla I
 946 Nco I
 972 Sac I
 998 BamH I
 1024 Xba I
 1050 Nco I
 1076 Mlu I
 1102 BamH I
 1128 Pst I
 1154 Cla I
 1180 EcoR V
 1206 Cla I
 1232 BamH I
 1258 Xba I
 1284 EcoR I
 1310 Not I
 1336 Sac I
 1362 BatX I
 1388 Sac I
 1414 Pvu I
 1440 Pvu I
 1466 Pvu I
 1492 Pvu I
 1518 Pvu I
 1544 Pvu I
 1570 Pvu I
 1596 Pvu I
 1622 Pvu I
 1648 Pvu I
 1674 Pvu I
 1700 Pvu I
 1726 Pvu I
 1752 Pvu I
 1778 Pvu I
 1804 Pvu I
 1830 Pvu I
 1856 Pvu I
 1882 Pvu I
 1908 Pvu I
 1934 Pvu I
 1960 Pvu I
 1986 Pvu I
 2012 Pvu I
 2038 Pvu I
 2064 Pvu I
 2090 Pvu I
 2116 Pvu I
 2142 Pvu I
 2168 Pvu I
 2194 Pvu I
 2220 Pvu I
 2246 Pvu I
 2272 Pvu I
 2298 Pvu I
 2324 Pvu I
 2350 Pvu I
 2376 Pvu I
 2402 Pvu I
 2428 Pvu I
 2454 Pvu I
 2480 Pvu I
 2506 Pvu I
 2532 Pvu I
 2558 Pvu I
 2584 Pvu I
 2610 Pvu I
 2636 Pvu I
 2662 Pvu I
 2688 Pvu I
 2714 Pvu I
 2740 Pvu I
 2766 Pvu I
 2792 Pvu I
 2818 Pvu I
 2844 Pvu I
 2870 Pvu I
 2896 Pvu I
 2922 Pvu I
 2948 Pvu I
 2974 Pvu I
 3000 Pvu I
 3026 Pvu I
 3052 Pvu I
 3078 Pvu I
 3104 Pvu I
 3130 Pvu I
 3156 Pvu I
 3182 Pvu I
 3208 Pvu I
 3234 Pvu I
 3260 Pvu I
 3286 Pvu I
 3312 Pvu I
 3338 Pvu I
 3364 Pvu I
 3390 Pvu I
 3416 Pvu I
 3442 Pvu I
 3468 Pvu I
 3494 Pvu I
 3520 Pvu I
 3546 Pvu I
 3572 Pvu I
 3598 Pvu I
 3624 Pvu I
 3650 Pvu I
 3676 Pvu I
 3702 Pvu I
 3728 Pvu I
 3754 Pvu I
 3780 Pvu I
 3806 Pvu I
 3832 Pvu I
 3858 Pvu I
 3884 Pvu I
 3910 Pvu I
 3936 Pvu I
 3962 Pvu I
 3988 Pvu I
 4014 Pvu I
 4040 Pvu I
 4066 Pvu I
 4092 Pvu I
 4118 Pvu I
 4144 Pvu I
 4170 Pvu I
 4196 Pvu I
 4222 Pvu I
 4248 Pvu I
 4274 Pvu I
 4300 Pvu I
 4326 Pvu I
 4352 Pvu I
 4378 Pvu I
 4404 Pvu I
 4430 Pvu I
 4456 Pvu I
 4482 Pvu I
 4508 Pvu I
 4534 Pvu I
 4560 Pvu I
 4586 Pvu I
 4612 Pvu I
 4638 Pvu I
 4664 Pvu I
 4690 Pvu I
 4716 Pvu I
 4742 Pvu I
 4768 Pvu I
 4794 Pvu I
 4820 Pvu I
 4846 Pvu I
 4872 Pvu I
 4898 Pvu I
 4924 Pvu I
 4950 Pvu I
 4976 Pvu I
 5002 Pvu I
 5028 Pvu I
 5054 Pvu I
 5080 Pvu I
 5106 Pvu I
 5132 Pvu I
 5158 Pvu I
 5184 Pvu I
 5210 Pvu I
 5236 Pvu I
 5262 Pvu I
 5288 Pvu I
 5314 Pvu I
 5340 Pvu I
 5366 Pvu I
 5392 Pvu I
 5418 Pvu I
 5444 Pvu I
 5470 Pvu I
 5496 Pvu I
 5522 Pvu I
 5548 Pvu I
 5574 Pvu I
 5600 Pvu I
 5626 Pvu I
 5652 Pvu I
 5678 Pvu I
 5704 Pvu I
 5730 Pvu I
 5756 Pvu I
 5782 Pvu I
 5808 Pvu I
 5834 Pvu I
 5860 Pvu I
 5886 Pvu I
 5912 Pvu I
 5938 Pvu I
 5964 Pvu I
 5990 Pvu I
 6016 Pvu I
 6042 Pvu I
 6068 Pvu I
 6094 Pvu I
 6120 Pvu I
 6146 Pvu I
 6172 Pvu I
 6198 Pvu I
 6224 Pvu I
 6250 Pvu I
 6276 Pvu I
 6302 Pvu I
 6328 Pvu I
 6354 Pvu I
 6380 Pvu I
 6406 Pvu I
 6432 Pvu I
 6458 Pvu I
 6484 Pvu I
 6510 Pvu I
 6536 Pvu I
 6562 Pvu I
 6588 Pvu I
 6614 Pvu I
 6640 Pvu I
 6666 Pvu I
 6692 Pvu I
 6718 Pvu I
 6744 Pvu I
 6770 Pvu I
 6796 Pvu I
 6822 Pvu I
 6848 Pvu I
 6874 Pvu I
 6900 Pvu I
 6926 Pvu I
 6952 Pvu I
 6978 Pvu I
 7004 Pvu I
 7030 Pvu I
 7056 Pvu I
 7082 Pvu I
 7108 Pvu I
 7134 Pvu I
 7160 Pvu I
 7186 Pvu I
 7212 Pvu I
 7238 Pvu I
 7264 Pvu I
 7290 Pvu I
 7316 Pvu I
 7342 Pvu I
 7368 Pvu I
 7394 Pvu I
 7420 Pvu I
 7446 Pvu I
 7472 Pvu I
 7498 Pvu I
 7524 Pvu I
 7550 Pvu I
 7576 Pvu I
 7602 Pvu I
 7628 Pvu I
 7654 Pvu I
 7680 Pvu I
 7706 Pvu I
 7732 Pvu I
 7758 Pvu I
 7784 Pvu I
 7810 Pvu I
 7836 Pvu I
 7862 Pvu I
 7888 Pvu I
 7914 Pvu I
 7940 Pvu I
 7966 Pvu I
 7992 Pvu I
 8018 Pvu I
 8044 Pvu I
 8070 Pvu I
 8096 Pvu I
 8122 Pvu I
 8148 Pvu I
 8174 Pvu I
 8200 Pvu I
 8226 Pvu I
 8252 Pvu I
 8278 Pvu I
 8304 Pvu I
 8330 Pvu I
 8356 Pvu I
 8382 Pvu I
 8408 Pvu I
 8434 Pvu I
 8460 Pvu I
 8486 Pvu I
 8512 Pvu I
 8538 Pvu I
 8564 Pvu I
 8590 Pvu I
 8616 Pvu I
 8642 Pvu I
 8668 Pvu I
 8694 Pvu I
 8720 Pvu I
 8746 Pvu I
 8772 Pvu I
 8798 Pvu I
 8824 Pvu I
 8850 Pvu I
 8876 Pvu I
 8902 Pvu I
 8928 Pvu I
 8954 Pvu I
 8980 Pvu I
 9006 Pvu I
 9032 Pvu I
 9058 Pvu I
 9084 Pvu I
 9110 Pvu I
 9136 Pvu I
 9162 Pvu I
 9188 Pvu I
 9214 Pvu I
 9240 Pvu I
 9266 Pvu I
 9292 Pvu I
 9318 Pvu I
 9344 Pvu I
 9370 Pvu I
 9396 Pvu I
 9422 Pvu I
 9448 Pvu I
 9474 Pvu I
 9500 Pvu I
 9526 Pvu I
 9552 Pvu I
 9578 Pvu I
 9604 Pvu I
 9630 Pvu I
 9656 Pvu I
 9682 Pvu I
 9708 Pvu I
 9734 Pvu I
 9760 Pvu I
 9786 Pvu I
 9812 Pvu I
 9838 Pvu I
 9864 Pvu I
 9890 Pvu I
 9916 Pvu I
 9942 Pvu I
 9968 Pvu I
 9994 Pvu I
 10020 Pvu I
 10046 Pvu I
 10072 Pvu I
 10098 Pvu I
 10124 Pvu I
 10150 Pvu I
 10176 Pvu I
 10202 Pvu I
 10228 Pvu I
 10254 Pvu I
 10280 Pvu I
 10306 Pvu I
 10332 Pvu I
 10358 Pvu I
 10384 Pvu I
 10410 Pvu I
 10436 Pvu I
 10462 Pvu I
 10488 Pvu I
 10514 Pvu I
 10540 Pvu I
 10566 Pvu I
 10592 Pvu I
 10618 Pvu I
 10644 Pvu I
 10670 Pvu I
 10696 Pvu I
 10722 Pvu I
 10748 Pvu I
 10774 Pvu I
 10800 Pvu I
 10826 Pvu I
 10852 Pvu I
 10878 Pvu I
 10904 Pvu I
 10930 Pvu I
 10956 Pvu I
 10982 Pvu I
 11008 Pvu I
 11034 Pvu I
 11060 Pvu I
 11086 Pvu I
 11112 Pvu I
 11138 Pvu I
 11164 Pvu I
 11190 Pvu I
 11216 Pvu I
 11242 Pvu I
 11268 Pvu I
 11294 Pvu I
 11320 Pvu I
 11346 Pvu I
 11372 Pvu I
 11398 Pvu I
 11424 Pvu I
 11450 Pvu I
 11476 Pvu I
 11502 Pvu I
 11528 Pvu I
 11554 Pvu I
 11580 Pvu I
 11606 Pvu I
 11632 Pvu I
 11658 Pvu I
 11684 Pvu I
 11710 Pvu I
 11736 Pvu I
 11762 Pvu I
 11788 Pvu I
 11814 Pvu I
 11840 Pvu I
 11866 Pvu I
 11892 Pvu I
 11918 Pvu I
 11944 Pvu I
 11970 Pvu I
 11996 Pvu I
 12022 Pvu I
 12048 Pvu I
 12074 Pvu I
 12100 Pvu I
 12126 Pvu I
 12152 Pvu I
 12178 Pvu I
 12204 Pvu I
 12230 Pvu I
 12256 Pvu I
 12282 Pvu I
 12308 Pvu I
 12334 Pvu I
 12360 Pvu I
 12386 Pvu I
 12412 Pvu I
 12438 Pvu I
 12464 Pvu I
 12490 Pvu I
 12516 Pvu I
 12542 Pvu I
 12568 Pvu I
 12594 Pvu I
 12620 Pvu I
 12646 Pvu I
 12672 Pvu I
 12698 Pvu I
 12724 Pvu I
 12750 Pvu I
 12776 Pvu I
 12802 Pvu I
 12828 Pvu I
 12854 Pvu I
 12880 Pvu I
 12906 Pvu I
 12932 Pvu I
 12958 Pvu I
 12984 Pvu I
 13010 Pvu I
 13036 Pvu I
 13062 Pvu I
 13088 Pvu I
 13114 Pvu I
 13140 Pvu I
 13166 Pvu I
 13192 Pvu I
 13218 Pvu I
 13244 Pvu I
 13270 Pvu I
 13296 Pvu I
 13322 Pvu I
 13348 Pvu I
 13374 Pvu I
 13400 Pvu I
 13426 Pvu I
 13452 Pvu I
 13478 Pvu I
 13504 Pvu I
 13530 Pvu I
 13556 Pvu I
 13582 Pvu I
 13608 Pvu I
 13634 Pvu I
 13660 Pvu I
 13686 Pvu I
 13712 Pvu I
 13738 Pvu I
 13764 Pvu I
 13790 Pvu I
 13816 Pvu I
 13842 Pvu I
 13868 Pvu I
 13894 Pvu I
 13920 Pvu I
 13946 Pvu I
 13972 Pvu I
 14000 Pvu I
 14026 Pvu I
 14052 Pvu I
 14078 Pvu I
 14104 Pvu I
 14130 Pvu I
 14156 Pvu I
 14182 Pvu I
 14208 Pvu I
 14234 Pvu I
 14260 Pvu I
 14286 Pvu I
 14312 Pvu I
 14338 Pvu I
 14364 Pvu I
 14390 Pvu I
 14416 Pvu I
 14442 Pvu I
 14468 Pvu I
 14494 Pvu I
 14520 Pvu I
 14546 Pvu I
 14572 Pvu I
 14598 Pvu I
 14624 Pvu I
 14650 Pvu I
 14676 Pvu I
 14702 Pvu I
 14728 Pvu I
 14754 Pvu I
 14780 Pvu I
 14806 Pvu I
 14832 Pvu I
 14858 Pvu I
 14884 Pvu I
 14910 Pvu I
 14936 Pvu I
 14962 Pvu I
 14988 Pvu I
 15014 Pvu I
 15040 Pvu I
 15066 Pvu I
 15092 Pvu I
 15118 Pvu I
 15144 Pvu I
 15170 Pvu I
 15196 Pvu I
 15222 Pvu I
 15248 Pvu I
 15274 Pvu I
 15300 Pvu I
 15326 Pvu I
 15352 Pvu I
 15378 Pvu I
 15404 Pvu I
 15430 Pvu I
 15456 Pvu I
 15482 Pvu I
 15508 Pvu I
 15534 Pvu I
 15560 Pvu I
 15586 Pvu I
 15612 Pvu I
 15638 Pvu I
 15664 Pvu I
 15690 Pvu I
 15716 Pvu I
 15742 Pvu I
 15768 Pvu I
 15794 Pvu I
 15820 Pvu I
 15846 Pvu I
 15872 Pvu I
 15898 Pvu I
 15924 Pvu I
 15950 Pvu I
 15976 Pvu I
 16002 Pvu I
 16028 Pvu I
 16054 Pvu I
 16080 Pvu I
 16106 Pvu I
 16132 Pvu I
 16158 Pvu I
 16184 Pvu I
 16210 Pvu I
 16236 Pvu I
 16262 Pvu I
 16288 Pvu I
 16314 Pvu I
 16340 Pvu I
 16366 Pvu I
 16392 Pvu I
 16418 Pvu I
 16444 Pvu I
 16470 Pvu I
 16496 Pvu I
 16522 Pvu I
 16548 Pvu I
 16574 Pvu I
 16600 Pvu I
 16626 Pvu I
 16652 Pvu I
 16678 Pvu I
 16704 Pvu I
 16730 Pvu I
 16756 Pvu I
 16782 Pvu I
 16808 Pvu I
 16834 Pvu I
 16860 Pvu I
 16886 Pvu I
 16912 Pvu I
 16938 Pvu I
 16964 Pvu I
 16990 Pvu I
 17016 Pvu I
 17042 Pvu I
 17068 Pvu I
 17094 Pvu I
 17120 Pvu I
 17146 Pvu I
 17172 Pvu I
 17198 Pvu I
 17224 Pvu I
 17250 Pvu I
 17276 Pvu I
 17302 Pvu I
 17328 Pvu I
 17354 Pvu I
 17380 Pvu I
 17406 Pvu I
 17432 Pvu I
 17458 Pvu I
 17484 Pvu I
 17510 Pvu I
 17536 Pvu I
 17562 Pvu I
 17588 Pvu I
 17614 Pvu I
 17640 Pvu I
 17666 Pvu I
 17692 Pvu I
 17718 Pvu I
 17744 Pvu I
 17770 Pvu I
 17796 Pvu I
 17822 Pvu I
 17848 Pvu I
 17874 Pvu I
 17900 Pvu I
 17926 Pvu I
 17952 Pvu I
 17978 Pvu I
 18004 Pvu I
 18030 Pvu I
 18056 Pvu I
 18082 Pvu I
 18108 Pvu I
 18134 Pvu I
 18160 Pvu I
 18186 Pvu I
 18212 Pvu I
 18238 Pvu I
 18264 Pvu I
 18290 Pvu I
 18316 Pvu I
 18342 Pvu I
 18368 Pvu I
 18394 Pvu I
 18420 Pvu I
 18446 Pvu I
 18472 Pvu I
 18498 Pvu I
 18524 Pvu I
 18550 Pvu I
 18576 Pvu I
 18602 Pvu I
 18628 Pvu I
 18654 Pvu I
 18680 Pvu I
 18706 Pvu I
 18732 Pvu I
 18758 Pvu I
 18784 Pvu I
 18810 Pvu I
 18836 Pvu I
 18862 Pvu I
 18888 Pvu I
 18914 Pvu I
 18940 Pvu I
 18966 Pvu I
 18992 Pvu I
 19018 Pvu I
 19044 Pvu I
 19070 Pvu I
 19096 Pvu I
 19122 Pvu I
 19148 Pvu I
 19174 Pvu I
 19200 Pvu I
 19226 Pvu I
 19252 Pvu I
 19278 Pvu I
 19304 Pvu I
 19330 Pvu I
 19356 Pvu I
 19382 Pvu I
 19408 Pvu I
 19434 Pvu I
 19460 Pvu I
 19486 Pvu I
 19512 Pvu I
 19538 Pvu I
 19564 Pvu I
 19590 Pvu I
 19616 Pvu I
 19642 Pvu I
 19668 Pvu I
 19694 Pvu I
 19720 Pvu I
 19746 Pvu I
 19772 Pvu I
 19798 Pvu I
 19824 Pvu I
 19850 Pvu I
 19876 Pvu I
 19902 Pvu I
 19928 Pvu I
 19954 Pvu I
 19980 Pvu I
 20006 Pvu I
 20032 Pvu I
 20058 Pvu I
 20084 Pvu I
 20110 Pvu I
 20136 Pvu I
 20162 Pvu I
 20188 Pvu I
 20214 Pvu I
 20240 Pvu I
 20266 Pvu I
 20292 Pvu I
 20318 Pvu I
 20344 Pvu I
 20370 Pvu I
 20396 Pvu I
 20422 Pvu I
 20448 Pvu I
 20474 Pvu I
 20500 Pvu I
 20526 Pvu I
 20552 Pvu I
 20578 Pvu I
 20604 Pvu I
 20630 Pvu I
 20656 Pvu I
 20682 Pvu I
 20708 Pvu I
 20734 Pvu I
 20760 Pvu I
 20786 Pvu I
 20812 Pvu I
 20838 Pvu I
 20864 Pvu I
 20890 Pvu I
 20916 Pvu I
 20942 Pvu I
 20968 Pvu I
 20994 Pvu I
 21020 Pvu I
 21046 Pvu I
 21072 Pvu I
 21098 Pvu I
 21124 Pvu I
 21150 Pvu I
 21176 Pvu I
 21202 Pvu I
 21228 Pvu I
 21254 Pvu I
 21280 Pvu I
 21306 Pvu I
 21332 Pvu I
 21358 Pvu I
 21384 Pvu I
 21410 Pvu I
 21436 Pvu I
 21462 Pvu I
 21488 Pvu I
 21514 Pvu I
 21540 Pvu I
 21566 Pvu I
 21592 Pvu I
 21618 Pvu I
 21644 Pvu I
 21670 Pvu I
 21696 Pvu I
 21722 Pvu I
 21748 Pvu I
 21774 Pvu I
 21800 Pvu I
 21826 Pvu I
 21852 Pvu I
 21878 Pvu I
 21904 Pvu I
 21930 Pvu I
 21956 Pvu I
 21982 Pvu I
 22008 Pvu I
 22034 Pvu I
 22060 Pvu I
 22086 Pvu I
 22112 Pvu I
 22138 Pvu I
 22164 Pvu I
 22190 Pvu I
 22216 Pvu I
 22242 Pvu I
 22268 Pvu I
 22294 Pvu I
 22320 Pvu I
 22346 Pvu I
 22372 Pvu I
 22398 Pvu I
 22424 Pvu I
 22450 Pvu I
 22476 Pvu I
 22502 Pvu I
 22528 Pvu I
 22554 Pvu I
 22580 Pvu I
 22606 Pvu I
 22632 Pvu I
 22658 Pvu I
 22684 Pvu I
 22710 Pvu I
 22736 Pvu I
 22762 Pvu I
 22788 Pvu I
 22814 Pvu I
 22840 Pvu I
 22866 Pvu I
 22892 Pvu I
 22918 Pvu I
 22944 Pvu I
 22970 Pvu I
 22996 Pvu I
 23022 Pvu I
 23048 Pvu I
 23074 Pvu I
 23100 Pvu I
 23126 Pvu I
 23152 Pvu I
 23178 Pvu I
 23204 Pvu I
 23230 Pvu I
 23256 Pvu I
 23282 Pvu I
 23308 Pvu I
 23334 Pvu I
 23360 Pvu I
 23386 Pvu I
 23412 Pvu I
 23438 Pvu I
 23464 Pvu I
 23490 Pvu I
 23516 Pvu I
 23542 Pvu I
 23568 Pvu I
 23594 Pvu I
 23620 Pvu I
 23646 Pvu I
 23672 Pvu I
 23698 Pvu I
 23724 Pvu I
 23750 Pvu I
 23776 Pvu I
 23802 Pvu I
 23828 Pvu I
 23854 Pvu I
 23880 Pvu I
 23906 Pvu I
 23932 Pvu I
 23958 Pvu I
 23984 Pvu I
 24010 Pvu I
 24036 Pvu I
 24062 Pvu I
 24088 Pvu I
 24114 Pvu I
 24140 Pvu I
 24166 Pvu I
 24192 Pvu I
 24218 Pvu I
 24244 Pvu I
 24270 Pvu I
 24296 Pvu I
 24322 Pvu I
 24348 Pvu I
 24374 Pvu I
 24400 Pvu I
 24426 Pvu I
 24452 Pvu I
 24478 Pvu I
 24504 Pvu I
 24530 Pvu I
 24556 Pvu I
 24582 Pvu I
 24608 Pvu I
 24634 Pvu I
 24660 Pvu I
 24686 Pvu I
 24712 Pvu I
 24738 Pvu I
 24764 Pvu I
 24790 Pvu I
 24816 Pvu I
 24842 Pvu I
 24868 Pvu I
 24894 Pvu I
 24920 Pvu I
 24946 Pvu I
 24972 Pvu I
 25000 Pvu I
 25026 Pvu I
 25052 Pvu I
 25078 Pvu I
 25104 Pvu I
 25130 Pvu I
 25156 Pvu I
 25182 Pvu I
 25208 Pvu I
 25234 Pvu I
 25260 Pvu I
 25286 Pvu I
 25312 Pvu I
 25338 Pvu I
 25364 Pvu I
 25390 Pvu I
 25416 Pvu I
 25442 Pvu I
 25468 Pvu I
 25494 Pvu I
 25520 Pvu I
 25546 Pvu I
 25572 Pvu I
 25598 Pvu I
 25624 Pvu I
 25650 Pvu I
 25676 Pvu I
 25702 Pvu I
 25728

5	Sac I
11	Kpn I
17	Apa I
22	Xho I
28	Sal I
39	Stu I
74	Xho I
75	Sac I
358	Spe I
605	Xba I
811	Nco I
806	Sma I
1047	BamH I
1065	Pst I
1094	Pvu II
1362	Pvu II
1492	Hind III
1499	Cla I
1517	Pst I
1519	EcoR I
1745	Bgl II
1785	Bgl II
2217	Bgl II
2235	Spe I
2269	Sph I
2273	EcoR V
2276	Xho I
2280	Bgl II
2466	Pvu II
3907	Pvu I




FIG. 7



9/11




FIG.9

→ Repeat CTACACC in tandem
 - - - - - Repeat TTTTA
 Repeat ATTCT

10/11





FIG.10

11/11



Promoter or Es. z gene
 Codified sequence for Esr2 gene
 Terminator Ca MV 35 S

FIG.11

Ref. _____

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

PLANT SEED ENDOSPERM-SPECIFIC PROMOTERS

the specification of which: *(check one)*

REGULAR OR DESIGN APPLICATION

- ☐ is attached hereto.
- ☐ was filed on _____ as application Serial No. _____ and was amended on _____ (if applicable).

PCT FILED APPLICATION ENTERING NATIONAL STAGE

- ☒ was described and claimed in International application No. PCT/FR0002596 filed on 19/09/2000 and as amended on _____ (if any).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

PRIORITY CLAIM

I hereby claim foreign priority benefits under 35 USC 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN APPLICATION(S)

Country	Application Number	Date of Filing (day, month, year)	Priority Claimed
FRANCE	9912305	01/10/99	YES

(Complete this part only if this is a continuing application.)

I hereby claim the benefit under 35 USC 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 USC 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT-international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status--patented, pending, abandoned)

POWER OF ATTORNEY

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from _____ as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

As a named inventor, I hereby appoint the registered patent attorneys represented by Customer No. 000466 to prosecute this application and transact all business in the Patent and Trademark Office connected therewith, including: Robert J. PATCH, Reg. No. 17,355, Andrew J. PATCH, Reg. No. 32,925, Robert F. HARGEST, Reg. No. 25,590, Benoît CASTEL, Reg. No. 35,041, Eric JENSEN, Reg. No. 37,855, Thomas W. PERKINS, Reg. No. 33,027, and Roland E. LONG, Jr., Reg. No. 41,949,

c/o YOUNG & THOMPSON,
Second Floor,
745 South 23rd Street,
Arlington, Virginia 22202.



00466
PATENT, TRADEMARK OFFICE

Address all telephone calls to Young & Thompson at 703/521-2297. Telefax: 703/685-0573.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor: Jean-François BONELLO
(given name, family name)

Inventor's signature [Signature] Date 25/03/02
Residence: 9 Rue J. Monet - 77600 BUSSY ST GEORGES - FRANCE Citizenship: French
FRX

Post Office Address: The same as above

Full name of second joint inventor, if any: Peter ROGOWSKY
(given name, family name)

Inventor's signature [Signature] Date 25/03/02
Residence: 31 Rue André Bollier - 69007 LYON - FRANCE Citizenship: German
FRX

Post Office Address: The same as above

Full name of third joint inventor, if any: Pascual PEREZ
(given name, family name)

Inventor's signature [Signature] Date 25/03/02
Residence: 17 Chemin de la Pradelle VARENNES - 63450 CHANONAT Citizenship: French
FRX

Post Office Address: The same as above

Full name of fourth joint inventor:
(given name, family name)

Inventor's signature _____ Date _____
Residence: _____ Citizenship: _____

LISTE DE SEQUENCES

<110> Biogemma

<120> Promoteurs spécifiques de l'albumen des graines de végétaux

<130> BFF 99/496ext

<140>

<141>

<160> 21

<170> PatentIn Ver.. 2.1

<210> 1

<211> 531

<212> ADN

<213> Zea mays

<400> 1

gatcattaag	gactaagcag	tctttttccc	tttcggcttg	catcatcttt	agtcttcac	60
actattataa	gccgaagcta	ttaccccttg	gctatagctt	cggtgttcat	ctttattatc	120
ttcggactat	gtcttcacct	tgtatacctt	tgtcttgggg	gaaaaccttc	atcctgaagc	180
cgaagctccc	tgtaataatt	catatcatgc	taaaaataaa	tggtcagtc	tgtttttgag	240
gaccttcgga	agaggaaggc	cccccaacaa	gacgattaac	tagtattgtc	tactgcatc	300
gttttttttg	cacttcacat	ataatgcctc	aatagcatac	ttcattttag	gaactttatt	360
aaaactgtct	taaaataggg	ccaagtcata	aattcattca	aagtgactct	tcattttcta	420
cttcctatct	ttggtggttt	tgtatatata	tatgttcacg	gttgagtgat	gttcctacac	480
cactacacca	cacgttagat	atatatacag	aaaatagctt	cactatctag	a	531

<210> 2

<211> 2493

<212> ADN

<213> Zea mays

<400> 2

aagcttttcc	ggtgatgaag	cacctgtaat	acttaacagc	atgctgaaaa	caaatagtta	60
gctgtgtttt	tgaggacctt	cggaagatga	aggcccccac	cacatcccat	gcacaaagtc	120
cccatgactt	gcaaaaaagc	aaattttatc	aaaattttct	ataaaacact	tgaaaaacatt	180
tctctttttg	aaaagtgtag	agcactagca	actgtctact	aaaaagggtc	ccaaattttct	240
gggtataaca	atcgcatggg	aaataacaca	aaggaaatcc	tactaagagc	agtaattttg	300
ctaaaaaat	agtgcagcatt	ttaatgtaat	agggaatagg	agcatgcaat	acttgtgttc	360
tttcagggtt	ttgatgtcct	caaaagtgtg	ccccctggg	gcagttgcaa	cactcaaaat	420
ctactcgtat	acataaagaa	acatgggcac	aaaataagaa	acaatactca	aattatgaaa	480
aagggtcaaa	tggtcctata	attattgtag	acatttttaga	atttatttta	gacccaaaacc	540
atttaaat	ggttttaaat	gagttagata	ttaatattta	ttcagtttat	agttattttg	600
gcattttatt	tacttaacta	taacttctag	ggttttataa	gtaaattttg	ggtccctagt	660
tggaactagc	tacgattgct	ggttgatttc	cataaaagtc	gaggttcctt	tagcaaaaaat	720
ccacggtgaa	caagggggag	ataggtgttg	accgatattc	ctaaattttg	atcgttggac	780
ggcacatgga	tgtctcagat	taaatggtgg	atgtgcaagc	gacgcgcaca	cgatggagga	840
atggcttcac	gacggtgggc	tactagagct	ggctacgtca	accaatggag	ggctcgggtca	900
agggtcaaat	ttgttgccaa	gccactgtgg	ctcacgatga	gtcgattgag	cacatatcaa	960
ggtcgagggt	caaccagagg	ggcaagatcg	atggtgcagt	ggtgttctcg	atggaagggg	1020
aaacttcggt	gagcaattca	agatttccta	tcatgtgacc	gggtcagggg	atgggcgcac	1080
ggggtttggt	accttctggt	gcacatcatg	ttgctgtatc	gatgtcaagg	gagcattagg	1140
gttcacgagt	cagcgatgac	gggcatgggt	ggacttgtgt	cacatgggtt	cgatcaacta	1200

WO 01/25439

PCT/FR00/02596

2

```

gggacgatag agctctatga agtttcacaa ctctctcaca ctctagggat catgggtgaca 1260
aaggtgggga ggacggggcg tctctagtga ggggtggaatg cagttctgtc acgtggggaat 1320
agtggcggca tcgcttgtaa tgaataaaaag gtgcttgggg ggctgggaag tgcaatatga 1380
gggaagtagt tgggtcgggg atgttccttt tataagggag caccattgat taatggaaga 1440
caatgacaca aaggggtggt cgacagttta aagctcgaat gctgctaggg gtgctcaagg 1500
ttaaagatc aggcatacagg gaggaaggc agggataaaa tttctttact ccagttgtgg 1560
ggtgatgggg acaagggtag tgctcaagca agggaggggc agttcagcgc agagatgcct 1620
gttgtgacac atgggggggg gggaaattgga ggttgggggt gaccaggtga cgttatggcg 1680
tgacccagag aagagaccca ctgatgggga aaaaagggtgc caacaggtgg ggaccaaggt 1740
gtcagtgact caccgtgaca tgttattgga aagttacgtc cggaatgggt tgggcctgag 1800
tgatctagge tggctcgggc actgtgctga tcttttaatt tctccattcc caatttaagt 1860
tgaattttta attcaaatca aatgactcca aatctctcca aaattaccaa aatatagaat 1920
atttagatga atatgttggg ggagtttggg ctccgctttt ggtagtatg tttgtataaa 1980
aataatttct ctcttttgt cacttccaat attgacttaa atttttatgt agcaatgcc 2040
acttttttta gtagtgtgcc acttatagca caaaaactat atccattttc taatagtcct 2100
tgaaatccac attctatttt tagccattct tcaaaattgg cacaaaacta ggaaaattta 2160
atacattctt gccataacat attctagtgc aaatgttaac tagattgctc aatattagca 2220
aacttctttt gtaagattca ttaatatgce tacattgcat acttttttag aagttcatca 2280
ataatgcctc attagcattc ttcaatttag gaacttgatt aaaaccgcct taaaatagca 2340
ccaagtgcag gatccattta aaggtgattc ttaatttctt acttcctatc tttgggtggc 2400
tatgtttata tatgtgtggg tgggtgaatg atgttctac accactacac cacacgttgg 2460
acatatatat ggaaaatagc ttcacagtct aga 2493

```

<210> 3

<211> 1708

<212> ADN

<213> Zea mays

<400> 3

```

aagcttagaa attttaaaaa aagccaggca agcgttggtg tgcaaagagc taaaaattag 60
gaagacaaga gaacacggca agaaagcatg ctaaatgtgc tcgcggtgcg ttcttattta 120
tacgtcfaat acgttgcaag tggtagggcc ccactgtgca ttgactattg ctattctagc 180
aaaggaagg tgtttttcgg accttcggct taaggccttc gtccatatcg caatctgaat 240
ttatcattct aacaaattaa tattgtgagg ggctactgtt gggggccttc gacttccgaa 300
ggtcctcaaa aactggttta acagtgtttc tggagtataa tgcataaaca ggtatcttcg 360
ggtttggatc agaactacaa catgaagagg cacaaagaac acgaagggtg gcgcagagcc 420
gaagctcacg tgtaggagag cttcggcacg acagcagaaa aagggaaaccg acttaaaagg 480
aaaggctatt cagacctcga tggatttcta taggtcatta gcaaatgtaa agggcatgaa 540
tgtaatttta catgggctgt gtccttgccat ataaatagat gaacagtact ctgactgt 600
tcacgctgac ttggcattcg ctttttgcat cacgcttgta cccttgcttt ccttcaaacc 660
gaaggtagat ctataatttg ttattgtgtt attgtggata tggtaatgca aataaaaaata 720
agttgatgat aatgtttata ttatttttcg tatttcatat atgaattctt cctcatcatt 780
tattgtgctt acgaagggtt ttcttcaaaa atctttgtcc ggaattcatt atatccgaag 840
ggaaataatg tctcgaaggga cgaaggactt tgatatttaa cacttttcat gttgccttgt 900
tcttgactct tagcatttga gaacaagtc ccaacagctc ctaagctctt ctttgaagaa 960
acaactacta gatgaagttt ctccaaaagt acgtccattg aatggagtaa agagtcattt 1020
gacctctcgg aataaaaatta aaatgagaat aagtaagaat aaaacacctc tattatcaa 1080
tctaggccat acaaacattg ggtattacta aaaaatagct aatgccatct ttcaacattt 1140
ggaagttaaa accaaccaat cctcactcat tcccaagaaa tattggatca tatttaacat 1200
tttgtgtcac ttacaaaaat ggcttaattct tttatgcggc aatgccaaacc ttttttagca 1260
gggtgccact tgtaacatga aaactataac tattttcaaa tagtaccttg aaattcgc 1320
tctattttta tgcatcttcc aaaattgaca caaattaaac taggagaatt caatacat 1380
ttgccataac atattctaatt gcaaatatta agtagattgc tcaacatcgg tacacatctt 1440
ttgcaagatt aattagattt gtctcactac attctttgtt ttagcagttc atcaataatg 1500
cctcaatagc atacttcatt ttaggaactt ttgaaaaatt gtcttaaaat agggccaagt 1560
cacaaatcca cttcaaagggt gactcttcat ttcttacttc ctatctttgc ttgtttttgt 1620
atatatatgt gtggatgggt gagtgatgtt cctacaccac tacaccacac cttagacaca 1680
tatatggaaa atagcttcac tgtctaga 1708

```

WO 01/25439

PCT/FR00/02596

3

<210> 4
 <211> 1232
 <212> ADN
 <213> Zea mays

<400> 4
 tagttcatgc aaaagtagtg agtggtttata cacctatgcc aatgaataac ctctaactac 60
 cctcatgtac tggcctgggtg tttgggtaaac atgaagcaat aatttctctt tatcaaaata 120
 cgggtctcaa gctgcatgtg aaaaataata aatatttttt tgaagtgaat gaaaaatgat 180
 aaatatgaaa cagtaaatct ttccgttgaa aaagtacatc tctattatta acgataacct 240
 atatatcaat ctacaatgcg ctcatctgca tctcgatgca tactttcatc attttatgaa 300
 tgtacttttaa tgataagaag gattagaatg ttcttggttt cctcttattc ttaccttttt 360
 caaaattatc agtttccaat gtctgaatat gcaatgcatt ataaacccta gtcagcatat 420
 atcaagtcca tatataatgc tatatgttta agaactgggtg ctgagtatgt ctactcaaca 480
 tatttttttag ctattggatc gagcagttta gtaaaggtaa actacattta tctatcttca 540
 agttgtattt tcccaccctt aaattatgaa agggagtaac gctccactcc aactggtgaa 600
 agggaacaaa tttgggtctcc ggactgattt cattgggtgg tctctatttt ttaaaacaac 660
 aaaaaaacat atttggtctc tgaaaattga taattaatta atcataaatt aggaaaaaaa 720
 ctatattgaaa ctagtatatg ttttcttcta aaattattgt ctgtctgttg gtgctctagt 780
 tatagagtta taacatgaaa actatagcca ttttcaaata gtgccttgaa attcattttt 840
 gtttagccat tcttcaaaat tgccaccaa ccaggagaat ttattaataa attcacgcca 900
 taacatattc tagtccaaat gtttaagtga ttgctcaata tcattatact tcttttggac 960
 gattcattag tactgccttg ttgcatactt tgtttttagca gttcatcaat aatgcttcac 1020
 tagcataatt catgttagga acttgattaa aactgcctta agaacatggc aaagtgataa 1080
 atccacttca aaggcaatta ttaatttctt acttctatc tttgggtggt tttgtatata 1140
 tgtgtgtggg tgggttgatg atgcacacat ttctctacac cactacatca caccttggac 1200
 atatatgtga aaaaatagct tcgcagtcta ga 1232

<210> 5
 <211> 499
 <212> ADN
 <213> Zea mays

<400> 5
 ttttgtcact tccaatattg acttaaatct ttatgtagca atgccaaact tttttagtag 60
 tgtgccactt atagcacaaa aactatatcc attttcta atgccttgaa atccacattc 120
 tatttttagc cattcttcaa aattggcaca aaactaggaa aatttaatac attcttgcca 180
 taacatattc tagtgcaaat gtttaactaga ttgctcaata ttagcaaaact tcttttgtaa 240
 gattcattaa tattgttaca ttgcatactt ttttagaagt tcatcaataa tgcctcatta 300
 gcatacttca ttttaggaac ttgattaaaa ccgccttaaa atagagccaa gtgacggatc 360
 catttaaagg tgattcttaa tttcttactt cctatctttg gtggcttatg tttatatatg 420
 tgtgggtggg tgaatgatgt tcctacacca ctacaccaca cgttggacat atatatggaa 480
 aatagcttca cagtctaga 499

<210> 6
 <211> 507
 <212> ADN
 <213> Zea mays

<400> 6
 ttgtgtcact tacaaaaatg gcttaatctt ttatgcggca atgccaaact tttttagcag 60
 ggtgccactt gtaacatgaa aactataact attttcaaat agtaccttga aattcgcat 120
 ctatttttat gcattcttca aaattgacac aaattaaact aggagaattc aatacattct 180
 tgccataaca tatttctaag caaatattaa gtgagattgct caacatcggt acacatcttt 240
 tggacgatta attagtattg tctcactaca ttctttgttt tagcagttca tcaataatgc 300

WO 01/25439

PCT/FR00/02596

4

```

ctcaatagca tacttcattt taggaacttt atgaaaattg tcttaaaata gggccaagtc 360
acaaatccac ttcaaagggtg actcttcatt tcttacttcc tatctttgct tgtttttgta 420
tatatatgtg tggatgggtg agtgatgttc ctacaccact acaccacacc ttagacacat 480
atatggaaaa tagcttcact gtctaga 507

```

<210> 7
 <211> 265
 <212> ADN
 <213> Zea mays

```

<400> 7
namgattmay tartattggy wcaytrcatw sntntntttr gmasttcato aataatgcct 60
cawtagcata cttcatttta ggaacttkat kaaaayygyt ttaaaatagr gccaagtsay 120
rratycantt yaaagntgay tcttmatttc ttacttccta tctttgstkg yttwngtwta 180
tatatrtgtk srtgggtgar tgatgttctt acaccactac accacacstt rgayayatat 240
ayrgaaaata gcttcacwrt ctaga 265

```

<210> 8
 <211> 28
 <212> ADN
 <213> Séquence artificielle

<220>
 <223> Description de la séquence
 artificielle:oligonucléotide

```

<400> 8
ggggtctaga ctgtgaagct attttcca 28

```

<210> 9
 <211> 30
 <212> ADN
 <213> Séquence artificielle

<220>
 <223> Description de la séquence
 artificielle:oligonucléotide

```

<400> 9
ggggaagctt tacattcttg ccataacata 30

```

<210> 10
 <211> 30
 <212> ADN
 <213> Séquence artificielle

<220>
 <223> Description de la séquence
 artificielle:oligonucléotide

```

<400> 10
ggggaagctt ttcataata atgcctcatt 30

```

<210> 11

<211> 30
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 11
ggggaagctt taatttctta cttcctatct 30

<210> 12
<211> 27
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 12
ggccagtcga caaagcggcc gcatgca 27

<210> 13
<211> 19
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 13
tcagctgttt cgccggcgt 19

<210> 14
<211> 14
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 14
tcgactgcag ccca 14

<210> 15
<211> 14
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 15
gacgtcgggt tcga

14

<210> 16
<211> 16
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 16
ctagacccga attcgc

16

<210> 17
<211> 16
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 17
tgggcttaag cgccgg

16

<210> 18
<211> 15
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 18
gatccactag tcccg

15

<210> 19
<211> 15
<212> ADN
<213> Séquence artificielle

<220>
<223> Description de la séquence
artificielle:oligonucléotide

<400> 19
aattcgggac tagtg

15

<210> 20
<211> 17

<212> ADN

<213> Séquence artificielle

<220>

<223> Description de la séquence
artificielle:oligonucléotide

<400> 20

aagctttttg cggccgc

17

<210> 21

<211> 25

<212> ADN

<213> Séquence artificielle

<220>

<223> Description de la séquence
artificielle:oligonucléotide

<400> 21

tcgagcggcc gcaaaaagct tagct

25